

(09.12.04)



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ



REC'D 09 DEC 2004

WIPO

PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., c, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

Dated

4 November 2004

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)






15DEC03 0859224-4 002093  
POL/7700 0.00-0328906.3 ACCOUNT CHA

The Patent Office

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



1. Your reference

PPD 70311/GB/P

2. Patent application number

(The Patent Office will fill in this part)

0328906.3

12 DEC 2003

3. Full name, address and postcode of the or of each applicant (underline all surnames)

SYNGENTA PARTICIPATIONS AG  
Intellectual Property Department  
Schwarzwaldallee 215  
4058 Basel  
SWITZERLAND

Patents ADP number (if you know it)

8029555001

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

John Richard WATERMAN  
Intellectual Property Department  
Syngenta Limited  
Jealott's Hill International Research Centre  
PO Box 3538  
Bracknell, Berkshire, RG42 6YA  
UNITED KINGDOM

Patents ADP number (if you know it)

8029563001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

YES (b)

## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.  
Do not count copies of the same document

Continuation sheets of this form 80  
Description 03  
Claim(s) 01  
Abstract 00  
Drawing(s) 00

SN

10. If you are also filing any of the following, state how many against each item.

Priority documents  
Translations of priority documents  
Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)  
Request for preliminary examination and search (*Patents Form 9/77*)  
Request for substantive examination (*Patents Form 10/77*)  
Any other documents (*please specify*)

11. I/We request the grant of a patent on the basis of this application.

SYNGENTA PARTICIPATIONS AG  
Signature  
Authorised Signatory M A Rudd Date 12/12/03

12. Name and daytime telephone number of person to contact in the United Kingdom  
Margaret Ann RUDD 01344 413673

### Warning

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

### Notes


- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

# CHEMICAL COMPOUNDS

The present invention relates to spiroindane derivatives, to processes for preparing them, to insecticidal, acaricidal, molluscicidal and nematocidal compositions comprising them and to methods of using them to combat and control insect, acarine, mollusc and nematode pests.

Spiroindanes with pharmaceutical properties are disclosed in for example WO9808835, WO 9825604, WO 9417045, US 5434158, GB1421208 and GB1423851. It has now surprisingly been found that certain spiroindanes have insecticidal properties.

The present invention therefore provides a compound of formula (I):



wherein X is O or NR<sup>11</sup> where R<sup>11</sup> is hydrogen, optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

Y is a single bond, C=O, C=S or S(O)<sub>m</sub> where m is 0, 1 or 2;

R<sup>1</sup> is hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted alkylcarbonyl, aminocarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted heterocycloxy, cyano, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, formyl, optionally substituted heterocycl, optionally substituted alkylthio, NO or NR<sup>13</sup>R<sup>14</sup> where R<sup>13</sup> and R<sup>14</sup> are independently hydrogen, COR<sup>15</sup>, optionally substituted alkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocycl or R<sup>13</sup> and R<sup>14</sup> together with the N

atom to which they are attached form a group  $-N=C(R^{16})-NR^{17}R^{18}$ ;  $R^{15}$  is H, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aryloxy optionally substituted heteroaryl, optionally substituted heteroaryloxy or  $NR^{19}R^{20}$ ;  $R^{16}$ ,  $R^{17}$  and  $R^{18}$  are each independently H or lower alkyl;  $R^{19}$  and  $R^{20}$  are  
5 independently optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

$R^2$  and  $R^3$  are independently hydrogen, halogen, cyano, optionally substituted alkyl, optionally substituted alkoxy or optionally substituted aryl;

each  $R^4$  is independently halogen, nitro, cyano, optionally substituted  $C_{1-8}$  alkyl,  
10 optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted  $C_{3-7}$  cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy,  
15 optionally substituted heteroaryloxy, optionally substituted alkylthio or  $R^{21}R^{22}N$  where  $R^{21}$  and  $R^{22}$  are, independently, hydrogen,  $C_{1-8}$  alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-6}$  alkenyl,  $C_{3-6}$  alkynyl,  $C_{3-7}$  cycloalkyl( $C_{1-4}$ )alkyl,  $C_{2-6}$  haloalkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxycarbonyl or  $R^{21}$  and  $R^{22}$  together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from  
20 O, N or S and which may be optionally substituted by one or two  $C_{1-6}$  alkyl groups, or 2 adjacent groups  $R^4$  together with the carbon atoms to which they are attached form a 4, 5, 6, or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2, 3 or 4;

each  $R_a$  is independently hydrogen, halogen, hydroxy, cyano, optionally substituted  
25  $C_{1-8}$  alkyl, optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted  $C_{3-7}$  cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy,  
30 optionally substituted heteroaryloxy, optionally substituted alkylthio, optionally substituted

arylthio or  $R^{23}R^{24}N$  where  $R^{23}$  and  $R^{24}$  are, independently, hydrogen,  $C_{1-8}$  alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-6}$  alkenyl,  $C_{3-6}$  alkynyl,  $C_{3-7}$  cycloalkyl( $C_{1-4}$ )alkyl,  $C_{2-6}$  haloalkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxycarbonyl or  $R^{23}$  and  $R^{24}$  together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or  
5 two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two  $C_{1-6}$  alkyl groups, or two Ra groups attached to the same carbon atom are =O or two Ra groups attached to adjacent carbon atoms form a bond, or two Ra groups together with the carbon atom to which they are attached form a three- to seven-membered ring, that may be saturated or unsaturated, and that may contain one or two hetero atoms selected from  
10 the group consisting of N, O and S, and which may be optionally substituted by one or two  $C_{1-6}$  alkyl groups; or two Ra groups together form a group  $-CH_2-$ ,  $-CH=CH-$  or  $-CH_2CH_2-$ ; p is 0, 1, 2, 3, 4, 5 or 6; q is 0, 1, 2, 3, 4, 5 or 6 provided that p+q is 1, 2, 3, 4, 5 or 6;

$R^8$  is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally  
15 substituted alkoxy, optionally substituted aryloxy, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl or optionally substituted alkenylcarbonyl; or salts or N-oxides thereof.

The compounds of formula (I) may exist in different geometric or optical isomers or tautomeric forms. This invention covers all such isomers and tautomers and mixtures thereof  
20 in all proportions as well as isotopic forms such as deuterated compounds.

Each alkyl moiety either alone or as part of a larger group (such as alkoxy, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl) is a straight or branched chain and is, for example, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl or neo-pentyl. The alkyl groups are suitably  
25  $C_1$  to  $C_{12}$  alkyl groups, but are preferably  $C_1$ - $C_{10}$ , more preferably  $C_1$ - $C_8$ , even more preferably preferably  $C_1$ - $C_6$  and most preferably  $C_1$ - $C_4$  alkyl groups.

When present, the optional substituents on an alkyl moiety (alone or as part of a larger group such as alkoxy, alkoxycarbonyl, alkylcarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl) include one or more of halogen, nitro, cyano, NCS-,  $C_{3-7}$  cycloalkyl  
30 (itself optionally substituted with  $C_{1-6}$  alkyl or halogen),  $C_{5-7}$  cycloalkenyl (itself optionally

substituted with C<sub>1-6</sub> alkyl or halogen), hydroxy, C<sub>1-10</sub> alkoxy, C<sub>1-10</sub> alkoxy(C<sub>1-10</sub>)alkoxy, tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkoxycarbonyl(C<sub>1-10</sub>)alkoxy, C<sub>1-10</sub> haloalkoxy, aryl(C<sub>1-4</sub>)-alkoxy (where the aryl group is optionally substituted), C<sub>3-7</sub> cycloalkyloxy (where the cycloalkyl group is optionally substituted with C<sub>1-6</sub> alkyl or halogen), C<sub>2-10</sub> alkenyloxy, C<sub>2-10</sub> alkynyloxy, SH, C<sub>1-10</sub> alkylthio, C<sub>1-10</sub> haloalkylthio, aryl(C<sub>1-4</sub>)alkylthio (where the aryl group is optionally substituted), C<sub>3-7</sub> cycloalkylthio (where the cycloalkyl group is optionally substituted with C<sub>1-6</sub> alkyl or halogen), tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkylthio, arylthio (where the aryl group is optionally substituted), C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> haloalkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> haloalkylsulfinyl, arylsulfonyl (where the aryl group may be optionally substituted), tri(C<sub>1-4</sub>)alkylsilyl, aryldi(C<sub>1-4</sub>)alkylsilyl, (C<sub>1-4</sub>)alkyldiarylsilyl, triarylsilyl, C<sub>1-10</sub> alkylcarbonyl, HO<sub>2</sub>C, C<sub>1-10</sub> alkoxycarbonyl, aminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, di(C<sub>1-6</sub> alkyl)aminocarbonyl, N-(C<sub>1-3</sub> alkyl)-N-(C<sub>1-3</sub> alkoxy)aminocarbonyl, C<sub>1-6</sub> alkylcarbonyloxy, arylcarbonyloxy (where the aryl group is optionally substituted), di(C<sub>1-6</sub>)alkylaminocarbonyloxy, oximes such as =NOalkyl, =NOhaloalkyl and =NOaryl (itself optionally substituted), aryl (itself optionally substituted), heteroaryl (itself optionally substituted), heterocyclyl (itself optionally substituted with C<sub>1-6</sub> alkyl or halogen), aryloxy (where the aryl group is optionally substituted), heteroaryloxy, (where the heteroaryl group is optionally substituted), heterocyclyloxy (where the heterocyclyl group is optionally substituted with C<sub>1-6</sub> alkyl or halogen), amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub>)alkylamino, C<sub>1-6</sub> alkylcarbonylamino, N-(C<sub>1-6</sub>)alkylcarbonyl-N-(C<sub>1-6</sub>)alkylamino, C<sub>2-6</sub> alkenylcarbonyl, C<sub>2-6</sub> alkynylcarbonyl, C<sub>3-6</sub> alkenyloxycarbonyl, C<sub>3-6</sub> alkynyloxycarbonyl, aryloxycarbonyl (where the aryl group is optionally substituted) and arylcarbonyl (where the aryl group is optionally substituted).

Alkenyl and alkynyl moieties can be in the form of straight or branched chains, and the alkenyl moieties, where appropriate, can be of either the (*E*)- or (*Z*)-configuration. Examples are vinyl, allyl and propargyl.

When present, the optional substituents on alkenyl or alkynyl include those optional substituents given above for an alkyl moiety.

In the context of this specification acyl is optionally substituted C<sub>1-6</sub> alkylcarbonyl (for example acetyl), optionally substituted C<sub>2-6</sub> alkenylcarbonyl, optionally substituted C<sub>2-6</sub>



alkynylcarbonyl, optionally substituted arylcarbonyl (for example benzoyl) or optionally substituted heteroarylcarbonyl.

Halogen is fluorine, chlorine, bromine or iodine.

Haloalkyl groups are alkyl groups which are substituted with one or more of the same  
5 or different halogen atoms and are, for example, CF<sub>3</sub>, CF<sub>2</sub>Cl, CF<sub>3</sub>CH<sub>2</sub> or CHF<sub>2</sub>CH<sub>2</sub>.

In the context of the present specification the terms "aryl" and "aromatic ring system" refer to ring systems which may be mono-, bi- or tricyclic. Examples of such rings include phenyl, naphthalenyl, anthracenyl, indenyl or phenanthrenyl. A preferred aryl group is phenyl. In addition, the terms "heteroaryl", "heteroaromatic ring" or "heteroaromatic ring  
10 system" refer to an aromatic ring system containing at least one heteroatom and consisting either of a single ring or of two or more fused rings. Preferably, single rings will contain up to three and bicyclic systems up to four heteroatoms which will preferably be chosen from nitrogen, oxygen and sulphur. Examples of such groups include furyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl,  
15 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, 1,2,3-triazinyl, 1,2,4-triazinyl, 1,3,5-triazinyl, benzofuryl, benzisofuryl, benzothienyl, benzisothienyl, indolyl, isoindolyl, indazolyl, benzothiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, 2,1,3-benzoxadiazole  
20 quinolinyl, isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, benzotriazinyl, purinyl, pteridinyl and indolizinyl. Preferred examples of heteroaromatic radicals include pyridyl, pyrimidyl, triazinyl, thienyl, furyl, oxazolyl, isoxazolyl, 2,1,3-benzoxadiazole and thiazolyl.

The terms heterocycle and heterocyclyl refer to a non-aromatic ring containing up to  
25 10 atoms including one or more (preferably one or two) heteroatoms selected from O, S and N. Examples of such rings include 1,3-dioxolane, tetrahydrofuran and morpholine.

When present, the optional substituents on heterocyclyl include C<sub>1-6</sub> alkyl and C<sub>1-6</sub> haloalkyl as well as those optional substituents given above for an alkyl moiety.

Cycloalkyl includes cyclopropyl, cyclopentyl and cyclohexyl.

30 Cycloalkenyl includes cyclopentenyl and cyclohexenyl.

When present, the optional substituents on cycloalkyl or cycloalkenyl include C<sub>1-3</sub> alkyl as well as those optional substituents given above for an alkyl moiety.

Carbocyclic rings include aryl, cycloalkyl and cycloalkenyl groups.

When present, the optional substituents on aryl or heteroaryl are selected

5 independently, from halogen, nitro, cyano, NCS-, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy-  
(C<sub>1-6</sub>)alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> haloalkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> cycloalkyl (itself optionally  
substituted with C<sub>1-6</sub> alkyl or halogen), C<sub>5-7</sub> cycloalkenyl (itself optionally substituted with  
C<sub>1-6</sub> alkyl or halogen), hydroxy, C<sub>1-10</sub> alkoxy, C<sub>1-10</sub> alkoxy(C<sub>1-10</sub>)alkoxy, tri(C<sub>1-4</sub>)alkyl-  
silyl(C<sub>1-6</sub>)alkoxy, C<sub>1-6</sub> alkoxycarbonyl(C<sub>1-10</sub>)alkoxy, C<sub>1-10</sub> haloalkoxy, aryl(C<sub>1-4</sub>)alkoxy  
10 (where the aryl group is optionally substituted with halogen or C<sub>1-6</sub> alkyl), C<sub>3-7</sub> cycloalkyloxy  
(where the cycloalkyl group is optionally substituted with C<sub>1-6</sub> alkyl or halogen), C<sub>2-10</sub>  
alkenyloxy, C<sub>2-10</sub> alkynyloxy, SH, C<sub>1-10</sub> alkylthio, C<sub>1-10</sub> haloalkylthio, aryl(C<sub>1-4</sub>)alkylthio C<sub>3-7</sub>  
cycloalkylthio (where the cycloalkyl group is optionally substituted with C<sub>1-6</sub> alkyl or  
halogen), tri(C<sub>1-4</sub>)-alkylsilyl(C<sub>1-6</sub>)alkylthio, arylthio, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub>  
15 haloalkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> haloalkylsulfinyl, arylsulfonyl, tri(C<sub>1-4</sub>)alkylsilyl,  
aryldi(C<sub>1-4</sub>)-alkylsilyl, (C<sub>1-4</sub>)alkyldiarylsilyl, triarylsilyl, C<sub>1-10</sub> alkylcarbonyl, HO<sub>2</sub>C, C<sub>1-10</sub>  
alkoxycarbonyl, aminocarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, di(C<sub>1-6</sub> alkyl)-aminocarbonyl, N-  
(C<sub>1-3</sub> alkyl)-N-(C<sub>1-3</sub> alkoxy)aminocarbonyl, C<sub>1-6</sub> alkylcarbonyloxy, arylcarbonyloxy,  
di(C<sub>1-6</sub>)alkylamino-carbonyloxy, aryl (itself optionally substituted with C<sub>1-6</sub> alkyl or halogen),  
20 heteroaryl (itself optionally substituted with C<sub>1-6</sub> alkyl or halogen), heterocyclyl (itself  
optionally substituted with C<sub>1-6</sub> alkyl or halogen), aryloxy (where the aryl group is optionally  
substituted with C<sub>1-6</sub> alkyl or halogen), heteroaryloxy (where the heteroaryl group is  
optionally substituted with C<sub>1-6</sub> alkyl or halogen), heterocyclyloxy (where the heterocyclyl  
group is optionally substituted with C<sub>1-6</sub> alkyl or halogen), amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub>)  
25 alkylamino, C<sub>1-6</sub> alkylcarbonylamino, N-(C<sub>1-6</sub>)alkylcarbonyl-N-(C<sub>1-6</sub>)alkylamino,  
arylcarbonyl, (where the aryl group is itself optionally substituted with halogen or C<sub>1-6</sub> alkyl)  
or two adjacent positions on an aryl or heteroaryl system may be cyclised to form a 5, 6 or 7  
membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen or C<sub>1-6</sub>  
alkyl. Further substituents for aryl or heteroaryl include aryl carbonyl amino (where the aryl  
30 group is substituted by C<sub>1-6</sub> alkyl or halogen), (C<sub>1-6</sub>)alkyloxycarbonylamino

(C<sub>1-6</sub>)alkyloxycarbonyl-N-(C<sub>1-6</sub>)alkylamino, aryloxycarbonylamino (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), aryloxycarbonyl-N-(C<sub>1-6</sub>)alkylamino, (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), arylsulphonylamino (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), arylsulphonyl-N-(C<sub>1-6</sub>)alkylamino (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), aryl-N-(C<sub>1-6</sub>)alkylamino (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), arylamino (where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen), heteroaryl amino (where the heteroaryl group is substituted by C<sub>1-6</sub> alkyl or halogen), heterocyclylamino (where the heterocyclyl group is substituted by C<sub>1-6</sub> alkyl or halogen), aminocarbonylamino, C<sub>1-6</sub> alkylaminocarbonyl amino, di(C<sub>1-6</sub>)alkylaminocarbonyl amino, arylaminocarbonyl amino where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen, aryl-N-(C<sub>1-6</sub>)alkylaminocarbonylamino where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen, C<sub>1-6</sub> alkylaminocarbonyl-N-(C<sub>1-6</sub>)alkyl amino, di(C<sub>1-6</sub>)alkylaminocarbonyl-N-(C<sub>1-6</sub>)alkyl amino, arylaminocarbonyl-N-(C<sub>1-6</sub>)alkyl amino where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen) and aryl-N-(C<sub>1-6</sub>)alkylaminocarbonyl-N-(C<sub>1-6</sub>)alkyl amino where the aryl group is substituted by C<sub>1-6</sub> alkyl or halogen).

For substituted phenyl moieties, heterocyclyl and heteroaryl groups it is preferred that one or more substituents are independently selected from halogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> haloalkylthio, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> haloalkylsulfinyl, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> haloalkylsulfonyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> haloalkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> cycloalkyl, nitro, cyano, CO<sub>2</sub>H, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkoxycarbonyl, R<sup>25</sup>R<sup>26</sup>N or R<sup>27</sup>R<sup>28</sup>NC(O); wherein R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup> and R<sup>28</sup> are, independently, hydrogen or C<sub>1-6</sub> alkyl. Further preferred substituents are aryl and heteroaryl groups.

Haloalkenyl groups are alkenyl groups which are substituted with one or more of the same or different halogen atoms.

It is to be understood that dialkylamino substituents include those where the dialkyl groups together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which is optionally substituted by one or two independently selected (C<sub>1-6</sub>)alkyl groups. When heterocyclic rings are formed by joining two groups on an N atom,

the resulting rings are suitably pyrrolidine, piperidine, thiomorpholine and morpholine each of which may be substituted by one or two independently selected (C<sub>1-6</sub>) alkyl groups.

Preferably the optional substituents on an alkyl moiety include one or more of halogen, nitro, cyano, HO<sub>2</sub>C, C<sub>1-10</sub> alkoxy (itself optionally substituted by C<sub>1-10</sub> alkoxy),  
5 aryl(C<sub>1-4</sub>)alkoxy, C<sub>1-10</sub> alkylthio, C<sub>1-10</sub> alkylcarbonyl, C<sub>1-10</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, di(C<sub>1-6</sub> alkyl)aminocarbonyl, (C<sub>1-6</sub>)alkylcarbonyloxy, optionally substituted phenyl, heteroaryl, aryloxy, arylcarbonyloxy, heteroaryloxy, heterocyclyl, heterocyclyloxy, C<sub>3-7</sub> cycloalkyl (itself optionally substituted with (C<sub>1-6</sub>)alkyl or halogen), C<sub>3-7</sub> cycloalkyloxy, C<sub>5-7</sub> cycloalkenyl, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, tri(C<sub>1-4</sub>)alkylsilyl,  
10 tri(C<sub>1-4</sub>)alkylsilyl(C<sub>1-6</sub>)alkoxy, aryldi(C<sub>1-4</sub>)alkylsilyl, (C<sub>1-4</sub>)alkyldiarylsilyl and triarylsilyl.

Preferably the optional substituents on alkenyl or alkynyl include one or more of halogen, aryl and C<sub>3-7</sub> cycloalkyl.

A preferred optional substituent for heterocyclyl is C<sub>1-6</sub> alkyl.

Preferably the optional substituents for cycloalkyl include halogen, cyano and C<sub>1-3</sub>  
15 alkyl.

Preferably the optional substituents for cycloalkenyl include C<sub>1-3</sub> alkyl, halogen and cyano.

X is preferably NR<sup>11</sup>.

R<sup>11</sup> is preferably hydrogen or C<sub>1-6</sub> alkyl, most preferably hydrogen.

20 Preferably Y is a single bond, C=O or S(O)<sub>m</sub> where m is 0, 1 or 2.

More preferably Y is a single bond, C=O or SO<sub>2</sub>.

Yet more preferably Y is a single bond or C=O.

Most preferably Y is C=O.

Preferably R<sup>1</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> cyanoalkyl, C<sub>1-6</sub> haloalkyl, C<sub>3-7</sub>  
25 cycloalkyl(C<sub>1-4</sub>)alkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, heteroaryl(C<sub>1-6</sub>)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic  
30 ring, itself optionally substituted with halogen), aryl(C<sub>1-6</sub>)alkyl (wherein the aryl group may

be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the aryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C<sub>1-6</sub> alkylcarbonylamino(C<sub>1-6</sub>)alkyl, aryl (which may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the aryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), heteroaryl (which may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, phenoxy (wherein the phenyl group is optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), heteroaryloxy (optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), heterocyclyloxy (optionally substituted by halo, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), cyano, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-6</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, heterocyclyl (optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> haloalkylthio or NR<sup>13</sup>R<sup>14</sup> where R<sup>13</sup> and R<sup>14</sup> are independently hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, phenyl (which may be optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino, dialkylamino or C<sub>1-4</sub> alkoxycarbonyl), phenyl (C<sub>1-6</sub>)alkyl (wherein the phenyl group may be optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino, dialkylamino, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), heteroaryl (C<sub>1-6</sub>)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub>

haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen) or heteroaryl (which may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy, C<sub>1-4</sub> alkoxycarbonyl C<sub>1-6</sub> alkylcarbonylamino, phenyloxycarbonylamino (wherein the phenyl group is optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), amino, C<sub>1-6</sub> alkylamino or phenylamino (wherein the phenyl group is optionally substituted halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino)).

More preferably R<sup>1</sup> is C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, heteroaryl(C<sub>1-3</sub>)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), phenyl(C<sub>1-3</sub>)alkyl (wherein the phenyl group may be optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino, dialkylamino, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), phenyl (which may be optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino, dialkylamino, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), heteroaryl (which may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen), C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>2-6</sub> alkenyl, heterocyclyl (optionally substituted by halo, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> haloalkylthio or NR<sup>13</sup>R<sup>14</sup> where R<sup>13</sup> and R<sup>14</sup> are independently hydrogen, C<sub>1-6</sub>

alkyl or C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>2-6</sub> alkylcarbonyl, phenylcarbonyl, (where the phenyl is optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), phenyl(C<sub>1-3</sub>)alkyl (wherein the phenyl group may be optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino, dialkylamino, C<sub>1-6</sub> alkylsulfonfyl, C<sub>1-6</sub> alkoxycarbonyl, or two adjacent positions on the phenyl ring may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen) or heteroaryl(C<sub>1-3</sub>)alkyl (wherein the heteroaryl group may be optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, C<sub>1-6</sub> alkylsulfonfyl, C<sub>1-6</sub> alkylsulfinyl, C<sub>1-6</sub> alkylthio, C<sub>1-6</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylcarbonylamino, arylcarbonyl, or two adjacent positions on the heteroaryl system may be cyclised to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring, itself optionally substituted with halogen).

Even more preferably R<sup>1</sup> is C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, heteroaryl(C<sub>1-3</sub>)alkyl (wherein the heteroaryl group may be optionally substituted by halo, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl and where the heteroaryl group is a thiazole, pyridine, pyrimidine, pyrazine or pyridazine ring), heteroaryl (optionally substituted by halo, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl and where the heteroaryl group is a pyridine, pyrimidine, 2,1,3-benzoxadiazole, pyrazine or pyridazine ring), C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkylamino or heteroaryl(C<sub>1-3</sub>)alkylamino (wherein the heteroaryl group may be optionally substituted by halo, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl and where the heteroaryl group is a thiazole, pyridine, pyrimidine, pyrazine or pyridazine ring).

Most preferably R<sup>1</sup> is pyridyl (optionally substituted by halo, C<sub>1-3</sub> alkyl or C<sub>1-3</sub> haloalkyl) especially halo-substituted pyridyl.

It is preferred that R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or cyano.

More preferably R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halogen, C<sub>1-2</sub> alkyl, C<sub>1-2</sub> haloalkyl, C<sub>1-2</sub> alkoxy, cyano.

Even more preferably R<sup>2</sup> and R<sup>3</sup> are independently hydrogen or C<sub>1-4</sub> alkyl.

Yet more preferably R<sup>2</sup> and R<sup>3</sup> are independently hydrogen or methyl.

Most preferably  $R^2$  and  $R^3$  are both hydrogen.

Preferably each  $R^4$  is independently halogen, cyano,  $C_{1-8}$  alkyl,  $C_{1-8}$  haloalkyl,  $C_{1-6}$  cyanoalkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{3-7}$  cycloalkyl( $C_{1-6}$ )alkyl,  $C_{5-6}$  cycloalkenyl( $C_{1-6}$ )alkyl,  $C_{3-6}$  alkenyloxy( $C_{1-6}$ )alkyl,  $C_{3-6}$  alkynyloxy( $C_{1-6}$ )alkyl, aryloxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  carboxyalkyl,  $C_{1-6}$  alkylcarbonyl( $C_{1-6}$ )alkyl,  $C_{2-6}$  alkenylcarbonyl( $C_{1-6}$ )alkyl,  $C_{2-6}$  alkynylcarbonyl( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxy carbonyl( $C_{1-6}$ )alkyl,  $C_{3-6}$  alkenyloxycarbonyl( $C_{1-6}$ )alkyl,  $C_{3-6}$  alkynyloxycarbonyl( $C_{1-6}$ )alkyl, aryloxycarbonyl( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkylthio( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkylsulfinyl( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkylsulfonyl( $C_{1-6}$ )alkyl, aminocarbonyl( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkylaminocarbonyl( $C_{1-6}$ )alkyl, di( $C_{1-6}$ )alkylaminocarbonyl( $C_{1-6}$ )alkyl, phenyl( $C_{1-4}$ )alkyl (wherein the phenyl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), heteroaryl( $C_{1-4}$ )alkyl (wherein the heteroaryl group is optionally substituted by halo, nitro, cyano,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkoxy or  $C_{1-6}$  haloalkoxy), heterocyclyl( $C_{1-4}$ )alkyl (wherein the heterocyclyl group is optionally substituted by halo, nitro, cyano,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkoxy or  $C_{1-6}$  haloalkoxy),  $C_{2-6}$  alkenyl, aminocarbonyl( $C_{2-6}$ )alkenyl,  $C_{1-6}$  alkylaminocarbonyl( $C_{2-6}$ )alkenyl, di( $C_{1-6}$ )alkylaminocarbonyl( $C_{2-6}$ )alkenyl, phenyl( $C_{2-4}$ )alkenyl, (wherein the phenyl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino),  $C_{2-6}$  alkynyl, trimethylsilyl( $C_{2-6}$ )alkynyl, aminocarbonyl( $C_{2-6}$ )alkynyl,  $C_{1-6}$  alkylaminocarbonyl( $C_{2-6}$ )alkynyl, di( $C_{1-6}$ )alkylaminocarbonyl( $C_{2-6}$ )alkynyl,  $C_{1-6}$  alkoxy carbonyl,  $C_{3-7}$  cycloalkyl,  $C_{3-7}$  halocycloalkyl,  $C_{3-7}$  cyanocycloalkyl,  $C_{1-3}$  alkyl( $C_{3-7}$ )cycloalkyl,  $C_{1-3}$  alkyl( $C_{3-7}$ )halocycloalkyl, phenyl (optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), heteroaryl (optionally substituted by halo, nitro, cyano,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkoxy or  $C_{1-6}$  haloalkoxy), heterocyclyl (wherein the heterocyclyl group is optionally substituted by halo, nitro, cyano,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkoxy or  $C_{1-6}$  haloalkoxy), or 2 adjacent groups  $R^4$  together with the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen,  $C_{1-8}$  alkoxy,  $C_{1-6}$  haloalkoxy, phenoxy (optionally substituted by halo, nitro, cyano,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkoxy or  $C_{1-6}$  haloalkoxy),



heteroaryloxy (optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), C<sub>1-8</sub> alkylthio or R<sup>19</sup>R<sup>20</sup>N where R<sup>19</sup> and R<sup>20</sup> are, independently, hydrogen, C<sub>1-8</sub> alkyl, C<sub>3-7</sub> cycloalkyl, C<sub>3-6</sub> alkenyl, C<sub>3-6</sub> alkynyl, C<sub>2-6</sub> haloalkyl, C<sub>1-6</sub> alkoxycarbonyl or R<sup>19</sup> and R<sup>20</sup> together with the N atom to which they are attached form a  
5 five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two C<sub>1-6</sub> alkyl groups; n is 0, 1, 2 or 3.

More preferably each R<sup>4</sup> is independently halogen, cyano, C<sub>1-8</sub> alkyl, C<sub>1-8</sub> haloalkyl, C<sub>1-8</sub> cyanoalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>2-6</sub> alkynyl, trimethylsilyl(C<sub>2-6</sub>)alkynyl, C<sub>1-6</sub>  
10 alkoxycarbonyl, C<sub>3-7</sub> cycloalkyl, C<sub>1-3</sub> alkyl (C<sub>3-7</sub>) cycloalkyl, phenyl (optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), heterocyclyl (optionally substituted by halo, nitro, cyano, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> haloalkoxy), C<sub>1-8</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, phenoxy (optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy,  
15 CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), heteroaryloxy (optionally substituted by halo, nitro, cyano, C<sub>1-3</sub> alkyl, C<sub>1-3</sub> haloalkyl, C<sub>1-3</sub> alkoxy or C<sub>1-3</sub> haloalkoxy), di(C<sub>1-8</sub>)alkylamino, or 2 adjacent groups R<sup>4</sup> together with the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2 or 3.

Even more preferably each R<sup>4</sup> is independently halogen, cyano, C<sub>1-8</sub> alkyl, C<sub>1-8</sub> haloalkyl, C<sub>1-8</sub> cyanoalkyl, C<sub>1-6</sub> alkoxy(C<sub>1-6</sub>)alkyl, C<sub>2-6</sub> alkynyl, heterocyclyl (optionally substituted by C<sub>1-6</sub> alkyl), C<sub>1-8</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, phenoxy (optionally substituted by halo, cyano, C<sub>1-3</sub> alkyl or C<sub>1-3</sub> haloalkyl), heteroaryloxy (optionally substituted by halo, cyano, C<sub>1-3</sub> alkyl or C<sub>1-3</sub> haloalkyl), di(C<sub>1-8</sub>)alkylamino or 2 adjacent groups R<sup>4</sup> together with  
25 the carbon atoms to which they are attached form a 4, 5, 6 or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2 or 3.

Yet more preferably each R<sup>4</sup> is independently fluoro, chloro, bromo, cyano, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> cyanoalkyl or C<sub>1-3</sub> alkoxy(C<sub>1-3</sub>)alkyl; n is 0, 1 or 2.

Most preferably each R<sup>4</sup> is independently fluoro, chloro, bromo, C<sub>1-4</sub> alkyl or C<sub>1-4</sub>  
30 haloalkyl; n is 1 or 2.

Preferably  $R^8$  is  $C_{1-10}$  alkyl,  $C_{1-10}$  haloalkyl, aryl( $C_{1-6}$ )alkyl (wherein the aryl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), heteroaryl( $C_{1-6}$ )alkyl (wherein the heteroaryl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), arylcarbonyl-( $C_{1-6}$ )alkyl (wherein the aryl group may be optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino and the alkyl group may be optionally substituted by aryl),  $C_{2-8}$  alkenyl,  $C_{2-8}$  haloalkenyl, aryl( $C_{2-6}$ )-alkenyl (wherein the aryl group is optionally substituted halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino,  $C_{1-6}$  alkoxy carbonyl, or two adjacent substituents can cyclise to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring), heteroaryl( $C_{2-6}$ )-alkenyl (wherein the heteroaryl group is optionally substituted halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino,  $C_{1-6}$  alkoxy carbonyl, or two adjacent substituents can cyclise to form a 5, 6 or 7 membered carbocyclic or heterocyclic ring),  $C_{2-6}$  alkynyl, phenyl( $C_{2-6}$ )alkynyl (wherein the phenyl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino),  $C_{3-7}$  cycloalkyl,  $C_{1-6}$  alkoxy carbonyl,  $C_{1-6}$  alkylcarbonyl,  $C_{1-6}$  haloalkylcarbonyl or aryl( $C_{2-6}$ )alkenylcarbonyl (wherein the aryl group may be optionally substituted halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), or  $-C(R^{51})(R^{52})-[CR^{53}=CR^{54}]_Z-R^{55}$  where  $Z$  is 1 or 2,  $R^{51}$  and  $R^{52}$  are each independently H, halo or  $C_{1-2}$  alkyl,  $R^{53}$  and  $R^{54}$  are each independently H, halogen,  $C_{1-4}$  alkyl or  $C_{1-4}$  haloalkyl and  $R^{55}$  is optionally substituted aryl or optionally substituted heteroaryl.

More preferably  $R^8$  is phenyl( $C_{1-4}$ )alkyl (wherein the phenyl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), heteroaryl( $C_{1-6}$ )alkyl (wherein the heteroaryl group is optionally substituted halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy, CN,  $NO_2$ , aryl, heteroaryl, amino or dialkylamino), phenyl( $C_{2-6}$ )alkenyl (wherein the phenyl group is optionally substituted by halogen,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{1-4}$  haloalkoxy,

CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino), heteroaryl(C<sub>2-6</sub>)alkenyl (wherein the heteroaryl group is optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino) or phenyl(C<sub>2-6</sub>)alkynyl (wherein the phenyl group is optionally substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino, or –  
 5 C(R<sup>51</sup>)(R<sup>52</sup>)-[CR<sup>53</sup>=CR<sup>54</sup>]<sub>z</sub>-R<sup>55</sup> where z is 1 or 2, R<sup>51</sup> and R<sup>52</sup> are each independently H, halo or C<sub>1-2</sub> alkyl, R<sup>53</sup> and R<sup>54</sup> are each independently H, halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> haloalkyl and R<sup>55</sup> is optionally substituted aryl or optionally substituted heteroaryl.

Most preferably R<sup>8</sup> is –C(R<sup>51</sup>)(R<sup>52</sup>)-[CR<sup>53</sup>=CR<sup>54</sup>]<sub>z</sub>-R<sup>55</sup> where z is 1 or 2, preferably 1,  
 10 R<sup>51</sup> and R<sup>52</sup> are each independently H, halo or C<sub>1-2</sub> alkyl, R<sup>53</sup> and R<sup>54</sup> are each independently H, halogen, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> haloalkyl and R<sup>55</sup> is phenyl substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino or heteroaryl substituted by halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino.

15 R<sup>51</sup> and R<sup>52</sup> are preferably hydrogen.

R<sup>53</sup> and R<sup>54</sup> are preferably hydrogen or halogen, especially hydrogen.

R<sup>55</sup> is preferably phenyl substituted with one to three substituents selected from halogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>1-4</sub> haloalkoxy, CN, NO<sub>2</sub>, aryl, heteroaryl, amino or dialkylamino.

20 Preferably each R<sub>a</sub> is independently hydrogen, halo, cyano, C<sub>1-3</sub> alkyl, hydroxy or two R<sub>a</sub> groups together with the carbon atom to which they are attached form a carbonyl group

More preferably each R<sub>a</sub> is independently hydrogen, fluoro, methyl, hydroxy or two R<sub>a</sub> groups together with the carbon atom to which they are attached form a carbonyl group

Most preferably each R<sub>a</sub> is hydrogen.


25 Preferably p is 1, 2 or 3 and q is 1, 2 or 3 and p+q is 2 or 3.

More preferably p is 1 or 2 and q is 2.

Most preferably p and q are both 2.

The compounds in Tables I to XXIV below illustrate the compounds of the invention.

Table I provides 782 compounds of formula Ia



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table 1

Compound No	$R^8$	$R^{4a}$	$R^{4b}$	$R^{4c}$	$R^{4d}$
I-1	4-chlorobenzyl	H	H	H	H
I-2	Cinnamyl	H	H	H	H
I-3	4-chlorocinnamyl	H	H	H	H
I-4	4-fluorocinnamyl	H	H	H	H
I-5	4-bromocinnamyl	H	H	H	H
I-6	4-trifluoromethylcinnamyl	H	H	H	H
I-7	4-trifluoromethoxycinnamyl	H	H	H	H
I-8	4-pentafluoroethoxycinnamyl	H	H	H	H
I-9	4-methoxycinnamyl	H	H	H	H
I-10	4-ethoxycinnamyl	H	H	H	H
I-11	4-cyanocinnamyl	H	H	H	H
I-12	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	H
I-13	3-(4-chlorophenyl)-but-2-enyl	H	H	H	H
I-14	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	H
I-15	3-chloro-4-fluoro-cinnamyl	H	H	H	H
I-16	3,5-dichloro-cinnamyl	H	H	H	H
I-17	5-phenyl-penta-2,4-dienyl	H	H	H	H

I-18	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	H	H
I-19	3-naphthalen-2-yl-allyl	H	H	H	H
I-20	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	H
I-21	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	H
I-22	3-pyridin-4-yl-allyl	H	H	H	H
I-23	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	H
I-24	4-chlorobenzyl	H	F	H	H
I-25	Cinnamyl	H	F	H	H
I-26	4-chlorocinnamyl	H	F	H	H
I-27	4-fluorocinnamyl	H	F	H	H
I-28	4-bromocinnamyl	H	F	H	H
I-29	4-trifluoromethylcinnamyl	H	F	H	H
I-30	4-trifluoromethoxycinnamyl	H	F	H	H
I-31	4-pentafluoroethoxycinnamyl	H	F	H	H
I-32	4-methoxycinnamyl	H	F	H	H
I-33	4-ethoxycinnamyl	H	F	H	H
I-34	4-cyanocinnamyl	H	F	H	H
I-35	3-(6-chloro-pyridin-3-yl)-allyl	H	F	H	H
I-36	3-(4-chlorophenyl)-but-2-enyl	H	F	H	H
I-37	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	H	H
I-38	3-chloro-4-fluoro-cinnamyl	H	F	H	H
I-39	3,5-dichloro-cinnamyl	H	F	H	H
I-40	5-phenyl-penta-2,4-dienyl	H	F	H	H
I-41	4-isopropoxyloxycarbonylamino-cinnamyl	H	F	H	H
I-42	3-naphthalen-2-yl-allyl	H	F	H	H
I-43	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	H	H
I-44	3-(5-chloro-pyridin-2-yl)-allyl	H	F	H	H
I-45	3-pyridin-4-yl-allyl	H	F	H	H
I-46	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	H	H

I-47	4-chlorobenzyl	H	Cl	H	H
I-48	Cinnamyl	H	Cl	H	H
I-49	4-chlorocinnamyl	H	Cl	H	H
I-50	4-fluorocinnamyl	H	Cl	H	H
I-51	4-bromocinnamyl	H	Cl	H	H
I-52	4-trifluoromethylcinnamyl	H	Cl	H	H
I-53	4-trifluoromethoxycinnamyl	H	Cl	H	H
I-54	4-pentafluoroethoxycinnamyl	H	Cl	H	H
I-55	4-methoxycinnamyl	H	Cl	H	H
I-56	4-ethoxycinnamyl	H	Cl	H	H
I-57	4-cyanocinnamyl	H	Cl	H	H
I-58	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	H	H
I-59	3-(4-chlorophenyl)-but-2-enyl	H	Cl	H	H
I-60	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	H	H
I-61	3-chloro-4-fluoro-cinnamyl	H	Cl	H	H
I-62	3,5-dichloro-cinnamyl	H	Cl	H	H
I-63	5-phenyl-penta-2,4-dienyl	H	Cl	H	H
I-64	4-isopropylloxycarbonylamino-cinnamyl	H	Cl	H	H
I-65	3-naphthalen-2-yl-allyl	H	Cl	H	H
I-66	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	H	H
I-67	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	H	H
I-68	3-pyridin-4-yl-allyl	H	Cl	H	H
I-69	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	H	H
I-70	4-chlorobenzyl	H	Br	H	H
I-71	Cinnamyl	H	Br	H	H
I-72	4-chlorocinnamyl	H	Br	H	H
I-73	4-fluorocinnamyl	H	Br	H	H
I-74	4-bromocinnamyl	H	Br	H	H
I-75	4-trifluoromethylcinnamyl	H	Br	H	H

I-76	4-trifluoromethoxycinnamyl	H	Br	H	H
I-77	4-pentafluoroethoxycinnamyl	H	Br	H	H
I-78	4-methoxycinnamyl	H	Br	H	H
I-79	4-ethoxycinnamyl	H	Br	H	H
I-80	4-cyanocinnamyl	H	Br	H	H
I-81	3-(6-chloro-pyridin-3-yl)-allyl	H	Br	H	H
I-82	3-(4-chlorophenyl)-but-2-enyl	H	Br	H	H
I-83	3-(4-chlorophenyl)-3-fluoro-allyl	H	Br	H	H
I-84	3-chloro-4-fluoro-cinnamyl	H	Br	H	H
I-85	3,5-dichloro-cinnamyl	H	Br	H	H
I-86	5-phenyl-penta-2,4-dienyl	H	Br	H	H
I-87	4-isopropoxyloxycarbonylamino-cinnamyl	H	Br	H	H
I-88	3-naphthalen-2-yl-allyl	H	Br	H	H
I-89	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Br	H	H
I-90	3-(5-chloro-pyridin-2-yl)-allyl	H	Br	H	H
I-91	3-pyridin-4-yl-allyl	H	Br	H	H
I-92	3-(2-Chloro-pyridin-4-yl)-allyl	H	Br	H	H
I-93	4-chlorobenzyl	H	CN	H	H
I-94	Cinnamyl	H	CN	H	H
I-95	4-chlorocinnamyl	H	CN	H	H
I-96	4-fluorocinnamyl	H	CN	H	H
I-97	4-bromocinnamyl	H	CN	H	H
I-98	4-trifluoromethylcinnamyl	H	CN	H	H
I-99	4-trifluoromethoxycinnamyl	H	CN	H	H
I-100	4-pentafluoroethoxycinnamyl	H	CN	H	H
I-101	4-methoxycinnamyl	H	CN	H	H
I-102	4-ethoxycinnamyl	H	CN	H	H
I-103	4-cyanocinnamyl	H	CN	H	H
I-104	3-(6-chloro-pyridin-3-yl)-allyl	H	CN	H	H

I-105	3-(4-chlorophenyl)-but-2-enyl	H	CN	H	H
I-106	3-(4-chlorophenyl)-3-fluoro-allyl	H	CN	H	H
I-107	3-chloro-4-fluoro-cinnamyl	H	CN	H	H
I-108	3,5-dichloro-cinnamyl	H	CN	H	H
I-109	5-phenyl-penta-2,4-dienyl	H	CN	H	H
I-110	4-isopropoxyloxycarbonylamino-cinnamyl	H	CN	H	H
I-111	3-naphthalen-2-yl-allyl	H	CN	H	H
I-112	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CN	H	H
I-113	3-(5-chloro-pyridin-2-yl)-allyl	H	CN	H	H
I-114	3-pyridin-4-yl-allyl	H	CN	H	H
I-115	3-(2-Chloro-pyridin-4-yl)-allyl	H	CN	H	H
I-116	4-chlorobenzyl	H	OMe	H	H
I-117	Cinnamyl	H	OMe	H	H
I-118	4-chlorocinnamyl	H	OMe	H	H
I-119	4-fluorocinnamyl	H	OMe	H	H
I-120	4-bromocinnamyl	H	OMe	H	H
I-121	4-trifluoromethylcinnamyl	H	OMe	H	H
I-122	4-trifluoromethoxycinnamyl	H	OMe	H	H
I-123	4-pentafluoroethoxycinnamyl	H	OMe	H	H
I-124	4-methoxycinnamyl	H	OMe	H	H
I-125	4-ethoxycinnamyl	H	OMe	H	H
I-126	4-cyanocinnamyl	H	OMe	H	H
I-127	3-(6-chloro-pyridin-3-yl)-allyl	H	OMe	H	H
I-128	3-(4-chlorophenyl)-but-2-enyl	H	OMe	H	H
I-129	3-(4-chlorophenyl)-3-fluoro-allyl	H	OMe	H	H
I-130	3-chloro-4-fluoro-cinnamyl	H	OMe	H	H
I-131	3,5-dichloro-cinnamyl	H	OMe	H	H
I-132	5-phenyl-penta-2,4-dienyl	H	OMe	H	H
I-133	4-isopropoxyloxycarbonylamino-cinnamyl	H	OMe	H	H



I-134	3-naphthalen-2-yl-allyl	H	OMe	H	H
I-135	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	OMe	H	H
I-136	3-(5-chloro-pyridin-2-yl)-allyl	H	OMe	H	H
I-137	3-pyridin-4-yl-allyl	H	OMe	H	H
I-138	3-(2-Chloro-pyridin-4-yl)-allyl	H	OMe	H	H
I-139	4-chlorobenzyl	H	OCF <sub>3</sub>	H	H
I-140	Cinnamyl	H	OCF <sub>3</sub>	H	H
I-141	4-chlorocinnamyl	H	OCF <sub>3</sub>	H	H
I-142	4-fluorocinnamyl	H	OCF <sub>3</sub>	H	H
I-143	4-bromocinnamyl	H	OCF <sub>3</sub>	H	H
I-144	4-trifluoromethylcinnamyl	H	OCF <sub>3</sub>	H	H
I-145	4-trifluoromethoxycinnamyl	H	OCF <sub>3</sub>	H	H
I-146	4-pentafluoroethoxycinnamyl	H	OCF <sub>3</sub>	H	H
I-147	4-methoxycinnamyl	H	OCF <sub>3</sub>	H	H
I-148	4-ethoxycinnamyl	H	OCF <sub>3</sub>	H	H
I-149	4-cyanocinnamyl	H	OCF <sub>3</sub>	H	H
I-150	3-(6-chloro-pyridin-3-yl)-allyl	H	OCF <sub>3</sub>	H	H
I-151	3-(4-chlorophenyl)-but-2-enyl	H	OCF <sub>3</sub>	H	H
I-152	3-(4-chlorophenyl)-3-fluoro-allyl	H	OCF <sub>3</sub>	H	H
I-153	3-chloro-4-fluoro-cinnamyl	H	OCF <sub>3</sub>	H	H
I-154	3,5-dichloro-cinnamyl	H	OCF <sub>3</sub>	H	H
I-155	5-phenyl-penta-2,4-dienyl	H	OCF <sub>3</sub>	H	H
I-156	4-isopropylloxycarbonylamino-cinnamyl	H	OCF <sub>3</sub>	H	H
I-157	3-naphthalen-2-yl-allyl	H	OCF <sub>3</sub>	H	H
I-158	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	OCF <sub>3</sub>	H	H
I-159	3-(5-chloro-pyridin-2-yl)-allyl	H	OCF <sub>3</sub>	H	H
I-160	3-pyridin-4-yl-allyl	H	OCF <sub>3</sub>	H	H
I-161	3-(2-Chloro-pyridin-4-yl)-allyl	H	OCF <sub>3</sub>	H	H
I-162	4-chlorobenzyl	H	CH <sub>3</sub>	H	H

I-163	Cinnamyl	H	CH <sub>3</sub>	H	H
I-164	4-chlorocinnamyl	H	CH <sub>3</sub>	H	H
I-165	4-fluorocinnamyl	H	CH <sub>3</sub>	H	H
I-166	4-bromocinnamyl	H	CH <sub>3</sub>	H	H
I-167	4-trifluoromethylcinnamyl	H	CH <sub>3</sub>	H	H
I-168	4-trifluoromethoxycinnamyl	H	CH <sub>3</sub>	H	H
I-169	4-pentafluoroethoxycinnamyl	H	CH <sub>3</sub>	H	H
I-170	4-methoxycinnamyl	H	CH <sub>3</sub>	H	H
I-171	4-ethoxycinnamyl	H	CH <sub>3</sub>	H	H
I-172	4-cyanocinnamyl	H	CH <sub>3</sub>	H	H
I-173	3-(6-chloro-pyridin-3-yl)-allyl	H	CH <sub>3</sub>	H	H
I-174	3-(4-chlorophenyl)-but-2-enyl	H	CH <sub>3</sub>	H	H
I-175	3-(4-chlorophenyl)-3-fluoro-allyl	H	CH <sub>3</sub>	H	H
I-176	3-chloro-4-fluoro-cinnamyl	H	CH <sub>3</sub>	H	H
I-177	3,5-dichloro-cinnamyl	H	CH <sub>3</sub>	H	H
I-178	5-phenyl-penta-2,4-dienyl	H	CH <sub>3</sub>	H	H
I-179	4-isopropylloxycarbonylamino-cinnamyl	H	CH <sub>3</sub>	H	H
I-180	3-naphthalen-2-yl-allyl	H	CH <sub>3</sub>	H	H
I-181	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CH <sub>3</sub>	H	H
I-182	3-(5-chloro-pyridin-2-yl)-allyl	H	CH <sub>3</sub>	H	H
I-183	3-pyridin-4-yl-allyl	H	CH <sub>3</sub>	H	H
I-184	3-(2-Chloro-pyridin-4-yl)-allyl	H	CH <sub>3</sub>	H	H
I-185	4-chlorobenzyl	H	CF <sub>3</sub>	H	H
I-186	Cinnamyl	H	CF <sub>3</sub>	H	H
I-187	4-chlorocinnamyl	H	CF <sub>3</sub>	H	H
I-188	4-fluorocinnamyl	H	CF <sub>3</sub>	H	H
I-189	4-bromocinnamyl	H	CF <sub>3</sub>	H	H
I-190	4-trifluoromethylcinnamyl	H	CF <sub>3</sub>	H	H
I-191	4-trifluoromethoxycinnamyl	H	CF <sub>3</sub>	H	H

I-192	4-pentafluoroethoxycinnamyl	H	CF <sub>3</sub>	H	H
I-193	4-methoxycinnamyl	H	CF <sub>3</sub>	H	H
I-194	4-ethoxycinnamyl	H	CF <sub>3</sub>	H	H
I-195	4-cyanocinnamyl	H	CF <sub>3</sub>	H	H
I-196	3-(6-chloro-pyridin-3-yl)-allyl	H	CF <sub>3</sub>	H	H
I-197	3-(4-chlorophenyl)-but-2-enyl	H	CF <sub>3</sub>	H	H
I-198	3-(4-chlorophenyl)-3-fluoro-allyl	H	CF <sub>3</sub>	H	H
I-199	3-chloro-4-fluoro-cinnamyl	H	CF <sub>3</sub>	H	H
I-200	3,5-dichloro-cinnamyl	H	CF <sub>3</sub>	H	H
I-201	5-phenyl-penta-2,4-dienyl	H	CF <sub>3</sub>	H	H
I-202	4-isopropoxyloxycarbonylamino-cinnamyl	H	CF <sub>3</sub>	H	H
I-203	3-naphthalen-2-yl-allyl	H	CF <sub>3</sub>	H	H
I-204	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	CF <sub>3</sub>	H	H
I-205	3-(5-chloro-pyridin-2-yl)-allyl	H	CF <sub>3</sub>	H	H
I-206	3-pyridin-4-yl-allyl	H	CF <sub>3</sub>	H	H
I-207	3-(2-Chloro-pyridin-4-yl)-allyl	H	CF <sub>3</sub>	H	H
I-208	4-chlorobenzyl	H	H	Cl	H
I-209	Cinnamyl	H	H	Cl	H
I-210	4-chlorocinnamyl	H	H	Cl	H
I-211	4-fluorocinnamyl	H	H	Cl	H
I-212	4-bromocinnamyl	H	H	Cl	H
I-213	4-trifluoromethylcinnamyl	H	H	Cl	H
I-214	4-trifluoromethoxycinnamyl	H	H	Cl	H
I-215	4-pentafluoroethoxycinnamyl	H	H	Cl	H
I-216	4-methoxycinnamyl	H	H	Cl	H
I-217	4-ethoxycinnamyl	H	H	Cl	H
I-218	4-cyanocinnamyl	H	H	Cl	H
I-219	3-(6-chloro-pyridin-3-yl)-allyl	H	H	Cl	H
I-220	3-(4-chlorophenyl)-but-2-enyl	H	H	Cl	H

I-221	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	Cl	H
I-222	3-chloro-4-fluoro-cinnamyl	H	H	Cl	H
I-223	3,5-dichloro-cinnamyl	H	H	Cl	H
I-224	5-phenyl-penta-2,4-dienyl	H	H	Cl	H
I-225	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	Cl	H
I-226	3-naphthalen-2-yl-allyl	H	H	Cl	H
I-227	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	Cl	H
I-228	3-(5-chloro-pyridin-2-yl)-allyl	H	H	Cl	H
I-229	3-pyridin-4-yl-allyl	H	H	Cl	H
I-230	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	Cl	H
I-231	4-chlorobenzyl	H	H	F	H
I-232	Cinnamyl	H	H	F	H
I-233	4-chlorocinnamyl	H	H	F	H
I-234	4-fluorocinnamyl	H	H	F	H
I-235	4-bromocinnamyl	H	H	F	H
I-236	4-trifluoromethylcinnamyl	H	H	F	H
I-237	4-trifluoromethoxycinnamyl	H	H	F	H
I-238	4-pentafluoroethoxycinnamyl	H	H	F	H
I-239	4-methoxycinnamyl	H	H	F	H
I-240	4-ethoxycinnamyl	H	H	F	H
I-241	4-cyanocinnamyl	H	H	F	H
I-242	3-(6-chloro-pyridin-3-yl)-allyl	H	H	F	H
I-243	3-(4-chlorophenyl)-but-2-enyl	H	H	F	H
I-244	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	F	H
I-245	3-chloro-4-fluoro-cinnamyl	H	H	F	H
I-246	3,5-dichloro-cinnamyl	H	H	F	H
I-247	5-phenyl-penta-2,4-dienyl	H	H	F	H
I-248	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	F	H
I-249	3-naphthalen-2-yl-allyl	H	H	F	H

I250	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	F	H
I-251	3-(5-chloro-pyridin-2-yl)-allyl	H	H	F	H
I-252	3-pyridin-4-yl-allyl	H	H	F	H
I-253	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	F	H
I-254	4-chlorobenzyl	H	H	Br	H
I-255	Cinnamyl	H	H	Br	H
I-256	4-chlorocinnamyl	H	H	Br	H
I-257	4-fluorocinnamyl	H	H	Br	H
I-258	4-bromocinnamyl	H	H	Br	H
I-259	4-trifluoromethylcinnamyl	H	H	Br	H
I-260	4-trifluoromethoxycinnamyl	H	H	Br	H
I-261	4-pentafluoroethoxycinnamyl	H	H	Br	H
I-262	4-methoxycinnamyl	H	H	Br	H
I-263	4-ethoxycinnamyl	H	H	Br	H
I-264	4-cyanocinnamyl	H	H	Br	H
I-265	3-(6-chloro-pyridin-3-yl)-allyl	H	H	Br	H
I-266	3-(4-chlorophenyl)-but-2-enyl	H	H	Br	H
I-267	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	Br	H
I-268	3-chloro-4-fluoro-cinnamyl	H	H	Br	H
I-269	3,5-dichloro-cinnamyl	H	H	Br	H
I-270	5-phenyl-penta-2,4-dienyl	H	H	Br	H
I-271	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	Br	H
I-272	3-naphthalen-2-yl-allyl	H	H	Br	H
I-273	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	Br	H
I-274	3-(5-chloro-pyridin-2-yl)-allyl	H	H	Br	H
I-275	3-pyridin-4-yl-allyl	H	H	Br	H
I-276	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	Br	H
I-277	4-chlorobenzyl	H	H	OCF <sub>3</sub>	H
I-278	Cinnamyl	H	H	OCF <sub>3</sub>	H

I-279	4-chlorocinnamyl	H	H	OCF <sub>3</sub>	H
I-280	4-fluorocinnamyl	H	H	OCF <sub>3</sub>	H
I-281	4-bromocinnamyl	H	H	OCF <sub>3</sub>	H
I-282	4-trifluoromethylcinnamyl	H	H	OCF <sub>3</sub>	H
I-283	4-trifluoromethoxycinnamyl	H	H	OCF <sub>3</sub>	H
I-284	4-pentafluoroethoxycinnamyl	H	H	OCF <sub>3</sub>	H
I-285	4-methoxycinnamyl	H	H	OCF <sub>3</sub>	H
I-286	4-ethoxycinnamyl	H	H	OCF <sub>3</sub>	H
I-287	4-cyanocinnamyl	H	H	OCF <sub>3</sub>	H
I-288	3-(6-chloro-pyridin-3-yl)-allyl	H	H	OCF <sub>3</sub>	H
I-289	3-(4-chlorophenyl)-but-2-enyl	H	H	OCF <sub>3</sub>	H
I-290	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	OCF <sub>3</sub>	H
I-291	3-chloro-4-fluoro-cinnamyl	H	H	OCF <sub>3</sub>	H
I-292	3,5-dichloro-cinnamyl	H	H	OCF <sub>3</sub>	H
I-293	5-phenyl-penta-2,4-dienyl	H	H	OCF <sub>3</sub>	H
I-294	4-isopropylloxycarbonylamino-cinnamyl	H	H	OCF <sub>3</sub>	H
I-295	3-naphthalen-2-yl-allyl	H	H	OCF <sub>3</sub>	H
I-296	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	OCF <sub>3</sub>	H
I-297	3-(5-chloro-pyridin-2-yl)-allyl	H	H	OCF <sub>3</sub>	H
I-298	3-pyridin-4-yl-allyl	H	H	OCF <sub>3</sub>	H
I-299	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	OCF <sub>3</sub>	H
I-300	4-chlorobenzyl	H	H	CH <sub>3</sub>	H
I-301	Cinnamyl	H	H	CH <sub>3</sub>	H
I-302	4-chlorocinnamyl	H	H	CH <sub>3</sub>	H
I-303	4-fluorocinnamyl	H	H	CH <sub>3</sub>	H
I-304	4-bromocinnamyl	H	H	CH <sub>3</sub>	H
I-305	4-trifluoromethylcinnamyl	H	H	CH <sub>3</sub>	H
I-306	4-trifluoromethoxycinnamyl	H	H	CH <sub>3</sub>	H
I-307	4-pentafluoroethoxycinnamyl	H	H	CH <sub>3</sub>	H

I-308	4-methoxycinnamyl	H	H	CH <sub>3</sub>	H
I-309	4-ethoxycinnamyl	H	H	CH <sub>3</sub>	H
I-310	4-cyanocinnamyl	H	H	CH <sub>3</sub>	H
I-311	3-(6-chloro-pyridin-3-yl)-allyl	H	H	CH <sub>3</sub>	H
I-312	3-(4-chlorophenyl)-but-2-enyl	H	H	CH <sub>3</sub>	H
I-313	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	CH <sub>3</sub>	H
I-314	3-chloro-4-fluoro-cinnamyl	H	H	CH <sub>3</sub>	H
I-315	3,5-dichloro-cinnamyl	H	H	CH <sub>3</sub>	H
I-316	5-phenyl-penta-2,4-dienyl	H	H	CH <sub>3</sub>	H
I-317	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	CH <sub>3</sub>	H
I-318	3-naphthalen-2-yl-allyl	H	H	CH <sub>3</sub>	H
I-319	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	CH <sub>3</sub>	H
I-320	3-(5-chloro-pyridin-2-yl)-allyl	H	H	CH <sub>3</sub>	H
I-321	3-pyridin-4-yl-allyl	H	H	CH <sub>3</sub>	H
I-322	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	CH <sub>3</sub>	H
I-323	4-chlorobenzyl	H	H	CF <sub>3</sub>	H
I-324	Cinnamyl	H	H	CF <sub>3</sub>	H
I-325	4-chlorocinnamyl	H	H	CF <sub>3</sub>	H
I-326	4-fluorocinnamyl	H	H	CF <sub>3</sub>	H
I-327	4-bromocinnamyl	H	H	CF <sub>3</sub>	H
I-328	4-trifluoromethylcinnamyl	H	H	CF <sub>3</sub>	H
I-329	4-trifluoromethoxycinnamyl	H	H	CF <sub>3</sub>	H
I-330	4-pentafluoroethoxycinnamyl	H	H	CF <sub>3</sub>	H
I-331	4-methoxycinnamyl	H	H	CF <sub>3</sub>	H
I-332	4-ethoxycinnamyl	H	H	CF <sub>3</sub>	H
I-333	4-cyanocinnamyl	H	H	CF <sub>3</sub>	H
I-334	3-(6-chloro-pyridin-3-yl)-allyl	H	H	CF <sub>3</sub>	H
I-335	3-(4-chlorophenyl)-but-2-enyl	H	H	CF <sub>3</sub>	H
I-336	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	CF <sub>3</sub>	H

I-337	3-chloro-4-fluoro-cinnamyl	H	H	CF <sub>3</sub>	H
I-338	3,5-dichloro-cinnamyl	H	H	CF <sub>3</sub>	H
I-339	5-phenyl-penta-2,4-dienyl	H	H	CF <sub>3</sub>	H
I-340	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	CF <sub>3</sub>	H
I-341	3-naphthalen-2-yl-allyl	H	H	CF <sub>3</sub>	H
I-342	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	CF <sub>3</sub>	H
I-343	3-(5-chloro-pyridin-2-yl)-allyl	H	H	CF <sub>3</sub>	H
I-344	3-pyridin-4-yl-allyl	H	H	CF <sub>3</sub>	H
I-345	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	CF <sub>3</sub>	H
I-346	4-chlorobenzyl	F	H	H	H
I-347	Cinnamyl	F	H	H	H
I-348	4-chlorocinnamyl	F	H	H	H
I-349	4-fluorocinnamyl	F	H	H	H
I-350	4-bromocinnamyl	F	H	H	H
I-351	4-trifluoromethylcinnamyl	F	H	H	H
I-352	4-trifluoromethoxycinnamyl	F	H	H	H
I-353	4-pentafluoroethoxycinnamyl	F	H	H	H
I-354	4-methoxycinnamyl	F	H	H	H
I-355	4-ethoxycinnamyl	F	H	H	H
I-356	4-cyanocinnamyl	F	H	H	H
I-357	3-(6-chloro-pyridin-3-yl)-allyl	F	H	H	H
I-358	3-(4-chlorophenyl)-but-2-enyl	F	H	H	H
I-359	3-(4-chlorophenyl)-3-fluoro-allyl	F	H	H	H
I-360	3-chloro-4-fluoro-cinnamyl	F	H	H	H
I-361	3,5-dichloro-cinnamyl	F	H	H	H
I-362	5-phenyl-penta-2,4-dienyl	F	H	H	H
I-363	4-isopropoxyloxycarbonylamino-cinnamyl	F	H	H	H
I-364	3-naphthalen-2-yl-allyl	F	H	H	H
I-365	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	H	H	H



I-366	3-(5-chloro-pyridin-2-yl)-allyl	F	H	H	H
I-367	3-pyridin-4-yl-allyl	F	H	H	H
I-368	3-(2-Chloro-pyridin-4-yl)-allyl	F	H	H	H
I-369	4-chlorobenzyl	Cl	H	H	H
I-370	Cinnamyl	Cl	H	H	H
I-371	4-chlorocinnamyl	Cl	H	H	H
I-372	4-fluorocinnamyl	Cl	H	H	H
I-373	4-bromocinnamyl	Cl	H	H	H
I-374	4-trifluoromethylcinnamyl	Cl	H	H	H
I-375	4-trifluoromethoxycinnamyl	Cl	H	H	H
I-376	4-pentafluoroethoxycinnamyl	Cl	H	H	H
I-377	4-methoxycinnamyl	Cl	H	H	H
I-378	4-ethoxycinnamyl	Cl	H	H	H
I-379	4-cyanocinnamyl	Cl	H	H	H
I-380	3-(6-chloro-pyridin-3-yl)-allyl	Cl	H	H	H
I-381	3-(4-chlorophenyl)-but-2-enyl	Cl	H	H	H
I-382	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	H	H	H
I-383	3-chloro-4-fluoro-cinnamyl	Cl	H	H	H
I-384	3,5-dichloro-cinnamyl	Cl	H	H	H
I-385	5-phenyl-penta-2,4-dienyl	Cl	H	H	H
I-386	4-isopropylloxycarbonylamino-cinnamyl	Cl	H	H	H
I-387	3-naphthalen-2-yl-allyl	Cl	H	H	H
I-388	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	H	H	H
I-389	3-(5-chloro-pyridin-2-yl)-allyl	Cl	H	H	H
I-390	3-pyridin-4-yl-allyl	Cl	H	H	H
I-391	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	H	H	H
I-392	4-chlorobenzyl	Br	H	H	H
I-393	Cinnamyl	Br	H	H	H
I-394	4-chlorocinnamyl	Br	H	H	H

I-395	4-fluorocinnamyl	Br	H	H	H
I-396	4-bromocinnamyl	Br	H	H	H
I-397	4-trifluoromethylcinnamyl	Br	H	H	H
I-398	4-trifluoromethoxycinnamyl	Br	H	H	H
I-399	4-pentafluoroethoxycinnamyl	Br	H	H	H
I-400	4-methoxycinnamyl	Br	H	H	H
I-401	4-ethoxycinnamyl	Br	H	H	H
I-402	4-cyanocinnamyl	Br	H	H	H
I-403	3-(6-chloro-pyridin-3-yl)-allyl	Br	H	H	H
I-404	3-(4-chlorophenyl)-but-2-enyl	Br	H	H	H
I-405	3-(4-chlorophenyl)-3-fluoro-allyl	Br	H	H	H
I-406	3-chloro-4-fluoro-cinnamyl	Br	H	H	H
I-407	3,5-dichloro-cinnamyl	Br	H	H	H
I-408	5-phenyl-penta-2,4-dienyl	Br	H	H	H
I-409	4-isopropoxyloxycarbonylamino-cinnamyl	Br	H	H	H
I-410	3-naphthalen-2-yl-allyl	Br	H	H	H
I-411	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Br	H	H	H
I-412	3-(5-chloro-pyridin-2-yl)-allyl	Br	H	H	H
I-413	3-pyridin-4-yl-allyl	Br	H	H	H
I-414	3-(2-Chloro-pyridin-4-yl)-allyl	Br	H	H	H
I-415	4-chlorobenzyl	CF <sub>3</sub>	H	H	H
I-416	Cinnamyl	CF <sub>3</sub>	H	H	H
I-417	4-chlorocinnamyl	CF <sub>3</sub>	H	H	H
I-418	4-fluorocinnamyl	CF <sub>3</sub>	H	H	H
I-419	4-bromocinnamyl	CF <sub>3</sub>	H	H	H
I-420	4-trifluoromethylcinnamyl	CF <sub>3</sub>	H	H	H
I-421	4-trifluoromethoxycinnamyl	CF <sub>3</sub>	H	H	H
I-422	4-pentafluoroethoxycinnamyl	CF <sub>3</sub>	H	H	H
I-423	4-methoxycinnamyl	CF <sub>3</sub>	H	H	H

I-424	4-ethoxycinnamyl	CF <sub>3</sub>	H	H	H
I-425	4-cyanocinnamyl	CF <sub>3</sub>	H	H	H
I-426	3-(6-chloro-pyridin-3-yl)-allyl	CF <sub>3</sub>	H	H	H
I-427	3-(4-chlorophenyl)-but-2-enyl	CF <sub>3</sub>	H	H	H
I-428	3-(4-chlorophenyl)-3-fluoro-allyl	CF <sub>3</sub>	H	H	H
I-429	3-chloro-4-fluoro-cinnamyl	CF <sub>3</sub>	H	H	H
I-430	3,5-dichloro-cinnamyl	CF <sub>3</sub>	H	H	H
I-431	5-phenyl-penta-2,4-dienyl	CF <sub>3</sub>	H	H	H
I-432	4-isopropylloxycarbonylamino-cinnamyl	CF <sub>3</sub>	H	H	H
I-433	3-naphthalen-2-yl-allyl	CF <sub>3</sub>	H	H	H
I-434	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	CF <sub>3</sub>	H	H	H
I-435	3-(5-chloro-pyridin-2-yl)-allyl	CF <sub>3</sub>	H	H	H
I-436	3-pyridin-4-yl-allyl	CF <sub>3</sub>	H	H	H
I-437	3-(2-Chloro-pyridin-4-yl)-allyl	CF <sub>3</sub>	H	H	H
I-438	4-chlorobenzyl	H	H	H	F
I-439	Cinnamyl	H	H	H	F
I-440	4-chlorocinnamyl	H	H	H	F
I-441	4-fluorocinnamyl	H	H	H	F
I-442	4-bromocinnamyl	H	H	H	F
I-443	4-trifluoromethylcinnamyl	H	H	H	F
I-444	4-trifluoromethoxycinnamyl	H	H	H	F
I-445	4-pentafluoroethoxycinnamyl	H	H	H	F
I-446	4-methoxycinnamyl	H	H	H	F
I-447	4-ethoxycinnamyl	H	H	H	F
I-448	4-cyanocinnamyl	H	H	H	F
I-449	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	F
I-450	3-(4-chlorophenyl)-but-2-enyl	H	H	H	F
I-451	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	F
I-452	3-chloro-4-fluoro-cinnamyl	H	H	H	F

I-453	3,5-dichloro-cinnamyl	H	H	H	F
I-454	5-phenyl-penta-2,4-dienyl	H	H	H	F
I-455	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	H	F
I-456	3-naphthalen-2-yl-allyl	H	H	H	F
I-457	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	F
I-458	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	F
I-459	3-pyridin-4-yl-allyl	H	H	H	F
I-460	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	F
I-461	4-chlorobenzyl	H	H	H	Cl
I-462	Cinnamyl	H	H	H	Cl
I-463	4-chlorocinnamyl	H	H	H	Cl
I-464	4-fluorocinnamyl	H	H	H	Cl
I-465	4-bromocinnamyl	H	H	H	Cl
I-466	4-trifluoromethylcinnamyl	H	H	H	Cl
I-467	4-trifluoromethoxycinnamyl	H	H	H	Cl
I-468	4-pentafluoroethoxycinnamyl	H	H	H	Cl
I-469	4-methoxycinnamyl	H	H	H	Cl
I-470	4-ethoxycinnamyl	H	H	H	Cl
I-471	4-cyanocinnamyl	H	H	H	Cl
I-472	3-(6-chloro-pyridin-3-yl)-allyl	H	H	H	Cl
I-473	3-(4-chlorophenyl)-but-2-enyl	H	H	H	Cl
I-474	3-(4-chlorophenyl)-3-fluoro-allyl	H	H	H	Cl
I-475	3-chloro-4-fluoro-cinnamyl	H	H	H	Cl
I-476	3,5-dichloro-cinnamyl	H	H	H	Cl
I-477	5-phenyl-penta-2,4-dienyl	H	H	H	Cl
I-478	4-isopropoxyloxycarbonylamino-cinnamyl	H	H	H	Cl
I-479	3-naphthalen-2-yl-allyl	H	H	H	Cl
I-480	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	H	H	Cl
I-481	3-(5-chloro-pyridin-2-yl)-allyl	H	H	H	Cl

I-482	3-pyridin-4-yl-allyl	H	H	H	Cl
I-483	3-(2-Chloro-pyridin-4-yl)-allyl	H	H	H	Cl
I-484	4-chlorobenzyl	H	F	F	H
I-485	Cinnamyl	H	F	F	H
I-486	4-chlorocinnamyl	H	F	F	H
I-487	4-fluorocinnamyl	H	F	F	H
I-488	4-bromocinnamyl	H	F	F	H
I-489	4-trifluoromethylcinnamyl	H	F	F	H
I-490	4-trifluoromethoxycinnamyl	H	F	F	H
I-491	4-pentafluoroethoxycinnamyl	H	F	F	H
I-492	4-methoxycinnamyl	H	F	F	H
I-493	4-ethoxycinnamyl	H	F	F	H
I-494	4-cyanocinnamyl	H	F	F	H
I-495	3-(6-chloro-pyridin-3-yl)-allyl	H	F	F	H
I-496	3-(4-chlorophenyl)-but-2-enyl	H	F	F	H
I-497	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	F	H
I-498	3-chloro-4-fluoro-cinnamyl	H	F	F	H
I-499	3,5-dichloro-cinnamyl	H	F	F	H
I-500	5-phenyl-penta-2,4-dienyl	H	F	F	H
I-501	4-isopropylloxycarbonylamino-cinnamyl	H	F	F	H
I-502	3-naphthalen-2-yl-allyl	H	F	F	H
I-503	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	F	H
I-504	3-(5-chloro-pyridin-2-yl)-allyl	H	F	F	H
I-505	3-pyridin-4-yl-allyl	H	F	F	H
I-506	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	F	H
I-507	4-chlorobenzyl	H	F	Cl	H
I-508	Cinnamyl	H	F	Cl	H
I-509	4-chlorocinnamyl	H	F	Cl	H
I-510	4-fluorocinnamyl	H	F	Cl	H

I-511	4-bromocinnamyl	H	F	Cl	H
I-512	4-trifluoromethylcinnamyl	H	F	Cl	H
I-513	4-trifluoromethoxycinnamyl	H	F	Cl	H
I-514	4-pentafluoroethoxycinnamyl	H	F	Cl	H
I-515	4-methoxycinnamyl	H	F	Cl	H
I-516	4-ethoxycinnamyl	H	F	Cl	H
I-517	4-cyanocinnamyl	H	F	Cl	H
I-518	3-(6-chloro-pyridin-3-yl)-allyl	H	F	Cl	H
I-519	3-(4-chlorophenyl)-but-2-enyl	H	F	Cl	H
I-520	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	Cl	H
I-521	3-chloro-4-fluoro-cinnamyl	H	F	Cl	H
I-522	3,5-dichloro-cinnamyl	H	F	Cl	H
I-523	5-phenyl-penta-2,4-dienyl	H	F	Cl	H
I-524	4-isopropylloxycarbonylamino-cinnamyl	H	F	Cl	H
I-525	3-naphthalen-2-yl-allyl	H	F	Cl	H
I-526	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	Cl	H
I-527	3-(5-chloro-pyridin-2-yl)-allyl	H	F	Cl	H
I-528	3-pyridin-4-yl-allyl	H	F	Cl	H
I-529	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	Cl	H
I-530	4-chlorobenzyl	H	Cl	F	H
I-531	Cinnamyl	H	Cl	F	H
I-532	4-chlorocinnamyl	H	Cl	F	H
I-533	4-fluorocinnamyl	H	Cl	F	H
I-534	4-bromocinnamyl	H	Cl	F	H
I-535	4-trifluoromethylcinnamyl	H	Cl	F	H
I-536	4-trifluoromethoxycinnamyl	H	Cl	F	H
I-537	4-pentafluoroethoxycinnamyl	H	Cl	F	H
I-538	4-methoxycinnamyl	H	Cl	F	H
I-539	4-ethoxycinnamyl	H	Cl	F	H

I-540	4-cyanocinnamyl	H	Cl	F	H
I-541	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	F	H
I-542	3-(4-chlorophenyl)-but-2-enyl	H	Cl	F	H
I-543	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	F	H
I-544	3-chloro-4-fluoro-cinnamyl	H	Cl	F	H
I-545	3,5-dichloro-cinnamyl	H	Cl	F	H
I-546	5-phenyl-penta-2,4-dienyl	H	Cl	F	H
I-547	4-isopropylloxycarbonylamino-cinnamyl	H	Cl	F	H
I-548	3-naphthalen-2-yl-allyl	H	Cl	F	H
I-549	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	F	H
I-550	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	F	H
I-551	3-pyridin-4-yl-allyl	H	Cl	F	H
I-552	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	F	H
I-553	4-chlorobenzyl	H	Cl	Cl	H
I-554	Cinnamyl	H	Cl	Cl	H
I-555	4-chlorocinnamyl	H	Cl	Cl	H
I-556	4-fluorocinnamyl	H	Cl	Cl	H
I-557	4-bromocinnamyl	H	Cl	Cl	H
I-558	4-trifluoromethylcinnamyl	H	Cl	Cl	H
I-559	4-trifluoromethoxycinnamyl	H	Cl	Cl	H
I-560	4-pentafluoroethoxycinnamyl	H	Cl	Cl	H
I-561	4-methoxycinnamyl	H	Cl	Cl	H
I-562	4-ethoxycinnamyl	H	Cl	Cl	H
I-563	4-cyanocinnamyl	H	Cl	Cl	H
I-564	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	Cl	H
I-565	3-(4-chlorophenyl)-but-2-enyl	H	Cl	Cl	H
I-566	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	Cl	H
I-567	3-chloro-4-fluoro-cinnamyl	H	Cl	Cl	H
I-568	3,5-dichloro-cinnamyl	H	Cl	Cl	H

I-569	5-phenyl-penta-2,4-dienyl	H	Cl	Cl	H
I-570	4-isopropoxyloxycarbonylamino-cinnamyl	H	Cl	Cl	H
I-571	3-naphthalen-2-yl-allyl	H	Cl	Cl	H
I-572	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	Cl	H
I-573	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	Cl	H
I-574	3-pyridin-4-yl-allyl	H	Cl	Cl	H
I-575	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	Cl	H
I-576	4-chlorobenzyl	H	-OCF <sub>2</sub> O-		H
I-577	Cinnamyl	H	-OCF <sub>2</sub> O-		H
I-578	4-chlorocinnamyl	H	-OCF <sub>2</sub> O-		H
I-579	4-fluorocinnamyl	H	-OCF <sub>2</sub> O-		H
I-580	4-bromocinnamyl	H	-OCF <sub>2</sub> O-		H
I-581	4-trifluoromethylcinnamyl	H	-OCF <sub>2</sub> O-		H
I-582	4-trifluoromethoxycinnamyl	H	-OCF <sub>2</sub> O-		H
I-583	4-pentafluoroethoxycinnamyl	H	-OCF <sub>2</sub> O-		H
I-584	4-methoxycinnamyl	H	-OCF <sub>2</sub> O-		H
I-585	4-ethoxycinnamyl	H	-OCF <sub>2</sub> O-		H
I-586	4-cyanocinnamyl	H	-OCF <sub>2</sub> O-		H
I-587	3-(6-chloro-pyridin-3-yl)-allyl	H	-OCF <sub>2</sub> O-		H
I-588	3-(4-chlorophenyl)-but-2-enyl	H	-OCF <sub>2</sub> O-		H
I-589	3-(4-chlorophenyl)-3-fluoro-allyl	H	-OCF <sub>2</sub> O-		H
I-590	3-chloro-4-fluoro-cinnamyl	H	-OCF <sub>2</sub> O-		H
I-591	3,5-dichloro-cinnamyl	H	-OCF <sub>2</sub> O-		H
I-592	5-phenyl-penta-2,4-dienyl	H	-OCF <sub>2</sub> O-		H
I-593	4-isopropoxyloxycarbonylamino-cinnamyl	H	-OCF <sub>2</sub> O-		H
I-594	3-naphthalen-2-yl-allyl	H	-OCF <sub>2</sub> O-		H
I-595	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	-OCF <sub>2</sub> O-		H
I-596	3-(5-chloro-pyridin-2-yl)-allyl	H	-OCF <sub>2</sub> O-		H
I-597	3-pyridin-4-yl-allyl	H	-OCF <sub>2</sub> O-		H



I-598	3-(2-Chloro-pyridin-4-yl)-allyl	H	-OCF <sub>2</sub> O-		H
I-599	4-chlorobenzyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-600	Cinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-601	4-chlorocinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-602	4-fluorocinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-603	4-bromocinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-604	4-trifluoromethylcinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-605	4-trifluoromethoxycinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-606	4-pentafluoroethoxycinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-607	4-methoxycinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-608	4-ethoxycinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-609	4-cyanocinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-610	3-(6-chloro-pyridin-3-yl)-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-611	3-(4-chlorophenyl)-but-2-enyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-612	3-(4-chlorophenyl)-3-fluoro-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-613	3-chloro-4-fluoro-cinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-614	3,5-dichloro-cinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-615	5-phenyl-penta-2,4-dienyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-616	4-isopropylloxycarbonylamino-cinnamyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-617	3-naphthalen-2-yl-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-618	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-619	3-(5-chloro-pyridin-2-yl)-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-620	3-pyridin-4-yl-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-621	3-(2-Chloro-pyridin-4-yl)-allyl	H	-C <sub>4</sub> H <sub>4</sub> -		H
I-622	4-chlorobenzyl	Cl	H	Cl	H
I-623	Cinnamyl	Cl	H	Cl	H
I-624	4-chlorocinnamyl	Cl	H	Cl	H
I-625	4-fluorocinnamyl	Cl	H	Cl	H
I-226	4-bromocinnamyl	Cl	H	Cl	H

I-627	4-trifluoromethylcinnamyl	Cl	H	Cl	H
I-628	4-trifluoromethoxycinnamyl	Cl	H	Cl	H
I-629	4-pentafluoroethoxycinnamyl	Cl	H	Cl	H
I-630	4-methoxycinnamyl	Cl	H	Cl	H
I-631	4-ethoxycinnamyl	Cl	H	Cl	H
I-632	4-cyanocinnamyl	Cl	H	Cl	H
I-633	3-(6-chloro-pyridin-3-yl)-allyl	Cl	H	Cl	H
I-634	3-(4-chlorophenyl)-but-2-enyl	Cl	H	Cl	H
I-635	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	H	Cl	H
I-636	3-chloro-4-fluoro-cinnamyl	Cl	H	Cl	H
I-637	3,5-dichloro-cinnamyl	Cl	H	Cl	H
I-638	5-phenyl-penta-2,4-dienyl	Cl	H	Cl	H
I-639	4-isopropyloxycarbonylamino-cinnamyl	Cl	H	Cl	H
I-640	3-naphthalen-2-yl-allyl	Cl	H	Cl	H
I-641	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	H	Cl	H
I-642	3-(5-chloro-pyridin-2-yl)-allyl	Cl	H	Cl	H
I-643	3-pyridin-4-yl-allyl	Cl	H	Cl	H
I-644	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	H	Cl	H
I-645	4-chlorobenzyl	Cl	Cl	H	H
I-646	Cinnamyl	Cl	Cl	H	H
I-647	4-chlorocinnamyl	Cl	Cl	H	H
I-648	4-fluorocinnamyl	Cl	Cl	H	H
I-649	4-bromocinnamyl	Cl	Cl	H	H
I-650	4-trifluoromethylcinnamyl	Cl	Cl	H	H
I-651	4-trifluoromethoxycinnamyl	Cl	Cl	H	H
I-652	4-pentafluoroethoxycinnamyl	Cl	Cl	H	H
I-653	4-methoxycinnamyl	Cl	Cl	H	H
I-654	4-ethoxycinnamyl	Cl	Cl	H	H
I-655	4-cyanocinnamyl	Cl	Cl	H	H

I-656	3-(6-chloro-pyridin-3-yl)-allyl	Cl	Cl	H	H
I-657	3-(4-chlorophenyl)-but-2-enyl	Cl	Cl	H	H
I-658	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	Cl	H	H
I-659	3-chloro-4-fluoro-cinnamyl	Cl	Cl	H	H
I-660	3,5-dichloro-cinnamyl	Cl	Cl	H	H
I-661	5-phenyl-penta-2,4-dienyl	Cl	Cl	H	H
I-662	4-isopropylloxycarbonylamino-cinnamyl	Cl	Cl	H	H
I-663	3-naphthalen-2-yl-allyl	Cl	Cl	H	H
I-664	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	Cl	H	H
I-665	3-(5-chloro-pyridin-2-yl)-allyl	Cl	Cl	H	H
I-666	3-pyridin-4-yl-allyl	Cl	Cl	H	H
I-667	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	Cl	H	H
I-668	4-chlorobenzyl	H	Cl	H	Cl
I-669	Cinnamyl	H	Cl	H	Cl
I-670	4-chlorocinnamyl	H	Cl	H	Cl
I-671	4-fluorocinnamyl	H	Cl	H	Cl
I-672	4-bromocinnamyl	H	Cl	H	Cl
I-673	4-trifluoromethylcinnamyl	H	Cl	H	Cl
I-674	4-trifluoromethoxycinnamyl	H	Cl	H	Cl
I-675	4-pentafluoroethoxycinnamyl	H	Cl	H	Cl
I-676	4-methoxycinnamyl	H	Cl	H	Cl
I-677	4-ethoxycinnamyl	H	Cl	H	Cl
I-678	4-cyanocinnamyl	H	Cl	H	Cl
I-679	3-(6-chloro-pyridin-3-yl)-allyl	H	Cl	H	Cl
I-680	3-(4-chlorophenyl)-but-2-enyl	H	Cl	H	Cl
I-681	3-(4-chlorophenyl)-3-fluoro-allyl	H	Cl	H	Cl
I-682	3-chloro-4-fluoro-cinnamyl	H	Cl	H	Cl
I-683	3,5-dichloro-cinnamyl	H	Cl	H	Cl
I-684	5-phenyl-penta-2,4-dienyl	H	Cl	H	Cl


I-685	4-isopropoxyloxycarbonylamino-cinnamyl	H	Cl	H	Cl
I-686	3-naphthalen-2-yl-allyl	H	Cl	H	Cl
I-687	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	Cl	H	Cl
I-688	3-(5-chloro-pyridin-2-yl)-allyl	H	Cl	H	Cl
I-689	3-pyridin-4-yl-allyl	H	Cl	H	Cl
I-690	3-(2-Chloro-pyridin-4-yl)-allyl	H	Cl	H	Cl
I-691	4-chlorobenzyl	H	F	H	F
I-692	Cinnamyl	H	F	H	F
I-693	4-chlorocinnamyl	H	F	H	F
I-694	4-fluorocinnamyl	H	F	H	F
I-695	4-bromocinnamyl	H	F	H	F
I-696	4-trifluoromethylcinnamyl	H	F	H	F
I-697	4-trifluoromethoxycinnamyl	H	F	H	F
I-698	4-pentafluoroethoxycinnamyl	H	F	H	F
I-699	4-methoxycinnamyl	H	F	H	F
I-700	4-ethoxycinnamyl	H	F	H	F
I-701	4-cyanocinnamyl	H	F	H	F
I-702	3-(6-chloro-pyridin-3-yl)-allyl	H	F	H	F
I-703	3-(4-chlorophenyl)-but-2-enyl	H	F	H	F
I-704	3-(4-chlorophenyl)-3-fluoro-allyl	H	F	H	F
I-705	3-chloro-4-fluoro-cinnamyl	H	F	H	F
I-706	3,5-dichloro-cinnamyl	H	F	H	F
I-707	5-phenyl-penta-2,4-dienyl	H	F	H	F
I-708	4-isopropoxyloxycarbonylamino-cinnamyl	H	F	H	F
I-709	3-naphthalen-2-yl-allyl	H	F	H	F
I-710	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	H	F	H	F
I-711	3-(5-chloro-pyridin-2-yl)-allyl	H	F	H	F
I-712	3-pyridin-4-yl-allyl	H	F	H	F
I-713	3-(2-Chloro-pyridin-4-yl)-allyl	H	F	H	F

I-714	4-chlorobenzyl	F	H	F	H
I-715	Cinnamyl	F	H	F	H
I-716	4-chlorocinnamyl	F	H	F	H
I-717	4-fluorocinnamyl	F	H	F	H
I-718	4-bromocinnamyl	F	H	F	H
I-719	4-trifluoromethylcinnamyl	F	H	F	H
I-720	4-trifluoromethoxycinnamyl	F	H	F	H
I-721	4-pentafluoroethoxycinnamyl	F	H	F	H
I-722	4-methoxycinnamyl	F	H	F	H
I-723	4-ethoxycinnamyl	F	H	F	H
I-724	4-cyanocinnamyl	F	H	F	H
I-725	3-(6-chloro-pyridin-3-yl)-allyl	F	H	F	H
I-726	3-(4-chlorophenyl)-but-2-enyl	F	H	F	H
I-727	3-(4-chlorophenyl)-3-fluoro-allyl	F	H	F	H
I-728	3-chloro-4-fluoro-cinnamyl	F	H	F	H
I-729	3,5-dichloro-cinnamyl	F	H	F	H
I-730	5-phenyl-penta-2,4-dienyl	F	H	F	H
I-731	4-isopropylloxycarbonylamino-cinnamyl	F	H	F	H
I-732	3-naphthalen-2-yl-allyl	F	H	F	H
I-733	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	H	F	H
I-734	3-(5-chloro-pyridin-2-yl)-allyl	F	H	F	H
I-735	3-pyridin-4-yl-allyl	F	H	F	H
I-736	3-(2-Chloro-pyridin-4-yl)-allyl	F	H	F	H
I-737	4-chlorobenzyl	F	F	H	H
I-738	Cinnamyl	F	F	H	H
I-739	4-chlorocinnamyl	F	F	H	H
I-740	4-fluorocinnamyl	F	F	H	H
I-741	4-bromocinnamyl	F	F	H	H
I-742	4-trifluoromethylcinnamyl	F	F	H	H

I-743	4-trifluoromethoxycinnamyl	F	F	H	H
I-744	4-pentafluoroethoxycinnamyl	F	F	H	H
I-745	4-methoxycinnamyl	F	F	H	H
I-746	4-ethoxycinnamyl	F	F	H	H
I-747	4-cyanocinnamyl	F	F	H	H
I-748	3-(6-chloro-pyridin-3-yl)-allyl	F	F	H	H
I-749	3-(4-chlorophenyl)-but-2-enyl	F	F	H	H
I-750	3-(4-chlorophenyl)-3-fluoro-allyl	F	F	H	H
I-751	3-chloro-4-fluoro-cinnamyl	F	F	H	H
I-752	3,5-dichloro-cinnamyl	F	F	H	H
I-753	5-phenyl-penta-2,4-dienyl	F	F	H	H
I-754	4-isopropylloxycarbonylamino-cinnamyl	F	F	H	H
I-755	3-naphthalen-2-yl-allyl	F	F	H	H
I-756	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	F	F	H	H
I-757	3-(5-chloro-pyridin-2-yl)-allyl	F	F	H	H
I-758	3-pyridin-4-yl-allyl	F	F	H	H
I-759	3-(2-Chloro-pyridin-4-yl)-allyl	F	F	H	H
I-760	4-chlorobenzyl	Cl	F	H	H
I-761	Cinnamyl	Cl	F	H	H
I-762	4-chlorocinnamyl	Cl	F	H	H
I-763	4-fluorocinnamyl	Cl	F	H	H
I-764	4-bromocinnamyl	Cl	F	H	H
I-765	4-trifluoromethylcinnamyl	Cl	F	H	H
I-766	4-trifluoromethoxycinnamyl	Cl	F	H	H
I-767	4-pentafluoroethoxycinnamyl	Cl	F	H	H
I-768	4-methoxycinnamyl	Cl	F	H	H
I-769	4-ethoxycinnamyl	Cl	F	H	H
I-770	4-cyanocinnamyl	Cl	F	H	H
I-771	3-(6-chloro-pyridin-3-yl)-allyl	Cl	F	H	H

I-772	3-(4-chlorophenyl)-but-2-enyl	Cl	F	H	H
I-773	3-(4-chlorophenyl)-3-fluoro-allyl	Cl	F	H	H
I-774	3-chloro-4-fluoro-cinnamyl	Cl	F	H	H
I-775	3,5-dichloro-cinnamyl	Cl	F	H	H
I-776	5-phenyl-penta-2,4-dienyl	Cl	F	H	H
I-777	4-isopropoxyloxycarbonylamino-cinnamyl	Cl	F	H	H
I-778	3-naphthalen-2-yl-allyl	Cl	F	H	H
I-779	3-(5-trifluoromethyl-pyridin-2-yl)-allyl	Cl	F	H	H
I-780	3-(5-chloro-pyridin-2-yl)-allyl	Cl	F	H	H
I-781	3-pyridin-4-yl-allyl	Cl	F	H	H
I-782	3-(2-Chloro-pyridin-4-yl)-allyl	Cl	F	H	H


Table II provides 782 compounds of formula Ib



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


5

Table III provides 782 compounds of formula Ic



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table IV provides 782 compounds of formula Id




5

wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table V provides 782 compounds of formula Ie





wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table VI provides 782 compounds of formula If



5


wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table VII provides 782 compounds of formula Ig



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table VIII provides 782 compounds of formula Ih



5


wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table IX provides 782 compounds of formula Ii



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table X provides 782 compounds of formula Ij



5


wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XI provides 782 compounds of formula Ik



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table XII provides 782 compounds of formula II



5

wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table XIII provides 782 compounds of formula Im



10

wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1


Table XIV provides 782 compounds of formula In



wherein the values of R<sup>8</sup>, R<sup>4a</sup>, R<sup>4b</sup>, R<sup>4c</sup> and R<sup>4d</sup> are given in Table 1

5


Table XV provides 782 compounds of formula Io



wherein the values of R<sup>8</sup>, R<sup>4a</sup>, R<sup>4b</sup>, R<sup>4c</sup> and R<sup>4d</sup> are given in Table 1


10

Table XVI provides 782 compounds of formula Ip




wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XVII provides 782 compounds of formula Iq




wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XVIII provides 782 compounds of formula Ir




10 wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XIX provides 782 compounds of formula Is




wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XX provides 782 compounds of formula It




wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XXI provides 782 compounds of formula Iu



wherein the values of  $R^8$ ,  $R^{4a}$ ,  $R^{4b}$ ,  $R^{4c}$  and  $R^{4d}$  are given in Table 1

Table XXII provides 15 compounds of formula Iv



wherein the values of  $R^{4b}$ ,  $R^{4c}$  and  $R^8$  are given in Table 2 together with LCMS/MS data obtained as for the data in Table 5 below.


Table 2

Compound	$R^{4b}$	$R^{4c}$	$R^8$	M.p.	LCMS (Ret. Time, min)	MS data
XXII.1	H	H	H	175-180	1'35	355



XXII.2	H	H	t-butoxycarbonyl	90-95	3'38	455
XXII.3	H	H	4-trifluoromethoxybenzyl		2'06	529
XXII.4	H	H	4-trifluoromethylbenzyl		2'01	513
XXII.5	H	H	4-isopropoxyloxycarbonylamino- benzyl		1'91	546
XXII.6	H	H	4-(2-Ethyl-2H-tetrazol-5-yl)- benzyl		1'88	541
XXII.7	H	H	4-cyanobenzyl		1'71	470
XXII.8	H	H	4-fluorobenzyl		1'74	463
XXII.9	H	H	benzyl		1'67	445
XXII.10	H	H	2,6-difluorobenzyl		1'70	481
XXII.11	H	H	3-chlorobenzyl		1'88	479
XXII.12	H	H	1-phenyl-ethyl		1'74	459
XXII.13	H	H	methyl		1'32	369
XXII.14	Cl	H	t-butoxycarbonyl	165	3'68	433, 473
XXII.15	H	Cl	t-butoxycarbonyl	165	3'65	433, 473

Table XXIII provides 23 compounds of formula Iw



wherein the values of Y and R<sup>1</sup> are given in Table 3 together with LCMS/MS data obtained as for the data in Table 5 below.


5

Table 3

Compound	Y	R <sup>1</sup>	M.p.	LCMS (Ret. Time, min)	LCM S (M+H)
XXIII.1	bond	H	140-142	1'70	366
XXIII.2	C=O	phenyl		2'00	470
XXIII.3	C=O	2-furyl		1'91	460
XXIII.4	C=O	3-pyridyl		1'82	471
XXIII.5	C=O	4-trifluoromethoxyphenylamino	225	2'65	569
XXIII.6	C=O	2,4-dichlorophenylamino	215-220	2'81	554
XXIII.7	C=O	4-methoxyphenylamino	270	2'38	515
XXIII.8	C=O	3-methoxyphenylamino	230	2'45	515
XXIII.9	C=O	2-chlorophenylamino		2'62	519
XXIII.10	C=O	3-chlorophenylamino		2'58	519
XXIII.11	C=O	4-chlorophenylamino	235	2'59	519
XXIII.12	C=O	3-trifluoromethyl-phenylamino	230	2'67	553
XXIII.13	C=O	4-trifluoromethylphenyl	200	2'37	538

XXIII.14	C=O	4-chlorophenyl	200	2'28	504
XXIII.15	C=O	phenylamino	230	2'35	485
XXIII.16	C=O	2-chlorophenyl	180	2'19	504
XXIII.17	C=O	2-hydroxyphenyl	230	2'10	486
XXIII.18	C=O	4-nitrophenyl	190	2'16	515
XXIII.19	C=O	3,5-di(trifluoromethyl)phenyl	190	2'60	506
XXIII.20	C=O	2-chloro-6-methoxy-pyrid-4-yl	210	2'31	535
XXIII.21	C=O	2-chloro-6-methyl-pyrid-4-yl	209	2'15	519
XXIII.22	C=O	2-amino-pyrid-3-yl	225	1'61	486
XXIII.23	C=O	4-trifluoromethyl-phenylamino		2'62	553

Table XXIV provides 3 compounds of formula Ix together with LCMS/MS data obtained as for the data in Table 5 below.



5 wherein the values of R<sup>1</sup> given in Table 4

Table 4

Compound	R <sup>1</sup>	M.p.	LCMS (Ret. Time, min)	MS data
XXIV.1	H	203	1'76	367/369
XXIV.2	CH <sub>3</sub>			

XXIV.3	benzyl			
--------	--------	--	--	--

Mass spectra data were obtained for selected compounds of Tables I to XXI using LCMS:

LC5: 254nm - gradient 10% A to 100% B A=H<sub>2</sub>O+0.01%HCOOH

5 B=CH<sub>3</sub>CN/CH<sub>3</sub>OH+0.01%HCOOH positive electrospray 150-1000 m/z.

The data are shown in Table 5.

Table 5


10

Compound	M.p.	LCMS (Ret. Time, min)	MS data
I.3		1'81	408
II.3		1'79	471
III.1	198-200	1'88	479
III.3	240	2'07	505
III.6	235	2'13	539
III.7	113	2'30	555
III.26	205	2'18	523
III.29	230	2'27	557
III.30	205	2'28	573
III.49	212	2'28	540
III.52		2'36	573
III.53		2'43	589
III.210	210	2'28	540
III.213		2'36	573
III.214		2'48	589
III.233	180	2'18	523

III.236	210	2'22	557
III.237		2'26	573
V.3	185	2'33	540
XII.3		1'84	424

The compounds of the invention may be made in a variety of ways. Thus for example they may be made by the reactions summarised in Scheme I.

## SCHEME I



Thus a compound of formula 1 may be synthesised from compounds of formula 2a by reaction with a compound of formula  $R_1YXNH_2$  where X is O or NH at a temperature of between ambient temperature and 120°C, in an organic solvent such as methanol, ethanol, isopropanol, 1,4-dioxan, benzene or toluene in the presence of an acid such as sulphuric acid  
5 or a base such as sodium hydroxide or sodium acetate.

A compound of formula 1 may also be synthesised from compounds of formula 3a by reaction with an alkylating agent of the formula  $R_8-L$ , where L is chloride, bromide, iodide or a sulfonate (e.g. mesylate or tosylate) or similar leaving group at a temperature of between ambient temperature and 100°C, typically ambient temperature, in an organic solvent such as  
10 acetonitrile, dimethylformamide, dichloromethane, chloroform or 1,2-dichloroethane in the presence of a tertiary amine base such as triethylamine or diisopropylethylamine and optionally catalysed by halide salts such as sodium iodide, potassium iodide or tetrabutylammonium iodide.

Alternatively, a compound of formula 3a may be reacted with an aldehyde of the  
15 formula  $RCHO$  at a temperature between ambient temperature and 100°C in an organic solvent such as tetrahydrofuran or ethanol or mixtures of solvents in the presence of a reducing agent such as borane-pyridine complex, sodium borohydride, sodium (triacetoxyl)borohydride, sodium cyanoborohydride or such like, to produce a compound of formula 1 where  $R_8$  is  $CH_2-R$ .

20 Similarly, a compound of formula 2a may be obtained from a compound of formula 2b by the methods described above for 3a.

A compound of formula 2b (or 3a) may be formed by reaction of a compound of formula 4 (or 3b) with an acid such as trifluoroacetic acid at ambient temperature in an organic solvent such as dichloromethane, chloroform or 1,2-dichloroethane followed by  
25 neutralisation with a base such as sodium bicarbonate.

A compound of formula 3b may be formed by reaction of a compound of formula 4 with a compound of formula  $R_1YXNH_2$  where X is O or NH at a temperature of between ambient temperature and 120°C, in an organic solvent such as methanol, ethanol, isopropanol, 1,4-dioxan, benzene or toluene in the presence of an acid such as sulphuric acid  
30 or a base such as sodium hydroxide or sodium acetate.


Compounds of formula 4 may be obtained from compounds of formula 5 by the methods described by Tata et al. in *Biorg. Med. Chem. Lett.* **1997**, 663-668.

Compounds of formula 5 may be obtained from indenenes of formula 6 by the methods described by Chambers et al. in *J. Med. Chem.* **1992**, 35, 2033-2039.

Certain compounds of formula 2a, 2b, 3a, 3b 4 and 5 are novel and as such form a further aspect of the invention.

5 Indenenes of formula 6 are either known compounds or may be prepared by known methods by a person skilled in the art. An example of those methods is given in *Bull. Soc. Chim. Fr.* **1973**, 11, 3092.

The skilled person will readily recognise that it is possible to interconvert one compound of formula I to other compounds of formula I and examples of such procedures  
10 are given in scheme II.





The compounds of formula (I) can be used to combat and control infestations of insect pests such as Lepidoptera, Diptera, Hemiptera, Thysanoptera, Orthoptera, Dictyoptera, Coleoptera, Siphonaptera, Hymenoptera and Isoptera and also other invertebrate pests, for example, acarine, nematode and mollusc pests. Insects, acarines, nematodes and molluscs are hereinafter collectively referred to as pests. The pests which may be combated and controlled by the use of the invention compounds include those pests associated with agriculture (which term includes the growing of crops for food and fibre products), horticulture and animal husbandry, companion animals, forestry and the storage of products of vegetable origin (such as fruit, grain and timber); those pests associated with the damage of man-made structures and the transmission of diseases of man and animals; and also nuisance pests (such as flies).

Examples of pest species which may be controlled by the compounds of formula (I) include: *Myzus persicae* (aphid), *Aphis gossypii* (aphid), *Aphis fabae* (aphid), *Lygus* spp. (capsids), *Dysdercus* spp. (capsids), *Nilaparvata lugens* (planthopper), *Nephotettix inciticeps* (leafhopper), *Nezara* spp. (stinkbugs), *Euschistus* spp. (stinkbugs), *Leptocorisa* spp. (stinkbugs), *Frankliniella occidentalis* (thrip), *Thrips* spp. (thrips), *Leptinotarsa decemlineata* (Colorado potato beetle), *Anthonomus grandis* (boll weevil), *Aonidiella* spp. (scale insects), *Trialeurodes* spp. (white flies), *Bemisia tabaci* (white fly), *Ostrinia nubilalis* (European corn borer), *Spodoptera littoralis* (cotton leafworm), *Heliothis virescens* (tobacco budworm), *Helicoverpa armigera* (cotton bollworm), *Helicoverpa zea* (cotton bollworm), *Sylepta derogata* (cotton leaf roller), *Pieris brassicae* (white butterfly), *Plutella xylostella* (diamond back moth), *Agrotis* spp. (cutworms), *Chilo suppressalis* (rice stem borer), *Locusta migratoria* (locust), *Chortiocetes terminifera* (locust), *Diabrotica* spp. (rootworms), *Panonychus ulmi* (European red mite), *Panonychus citri* (citrus red mite), *Tetranychus urticae* (two-spotted spider mite), *Tetranychus cinnabarinus* (carmine spider mite), *Phyllocoptruta oleivora* (citrus rust mite), *Polyphagotarsonemus latus* (broad mite), *Brevipalpus* spp. (flat mites), *Boophilus microplus* (cattle tick), *Dermacentor variabilis* (American dog tick), *Ctenocephalides felis* (cat flea), *Liriomyza* spp. (leafminer), *Musca domestica* (housefly), *Aedes aegypti* (mosquito), *Anopheles* spp. (mosquitoes), *Culex* spp. (mosquitoes), *Lucillia* spp. (blowflies), *Blattella germanica* (cockroach), *Periplaneta americana* (cockroach), *Blatta orientalis* (cockroach), termites of the Mastotermitidae (for example *Mastotermes* spp.), the Kalotermitidae (for example *Neotermes* spp.), the

Rhinotermitidae (for example *Coptotermes formosanus*, *Reticulitermes flavipes*, *R. speratu*, *R. virginicus*, *R. hesperus*, and *R. santonensis*) and the Termitidae (for example *Globitermes sulphureus*), *Solenopsis geminata* (fire ant), *Monomorium pharaonis* (pharaoh's ant), *Damalinia* spp. and *Linognathus* spp. (biting and sucking lice), *Meloidogyne* spp. (root knot  
5 nematodes), *Globodera* spp. and *Heterodera* spp. (cyst nematodes), *Pratylenchus* spp. (lesion nematodes), *Rhodopholus* spp. (banana burrowing nematodes), *Tylenchulus* spp. (citrus nematodes), *Haemonchus contortus* (barber pole worm), *Caenorhabditis elegans* (vinegar eelworm), *Trichostrongylus* spp. (gastro intestinal nematodes) and *Deroceras reticulatum* (slug).

10 The invention therefore provides a method of combating and controlling insects, acarines, nematodes or molluscs which comprises applying an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I), or a composition containing a compound of formula (I), to a pest, a locus of pest, or to a plant susceptible to attack by a pest, The compounds of formula (I) are preferably used against  
15 insects, acarines or nematodes.

The term "plant" as used herein includes seedlings, bushes and trees.

In order to apply a compound of formula (I) as an insecticide, acaricide, nematocide or molluscicide to a pest, a locus of pest, or to a plant susceptible to attack by a pest, a compound of formula (I) is usually formulated into a composition which includes, in addition  
20 to the compound of formula (I), a suitable inert diluent or carrier and, optionally, a surface active agent (SFA). SFAs are chemicals which are able to modify the properties of an interface (for example, liquid/solid, liquid/air or liquid/liquid interfaces) by lowering the interfacial tension and thereby leading to changes in other properties (for example dispersion, emulsification and wetting). It is preferred that all compositions (both solid and liquid  
25 formulations) comprise, by weight, 0.0001 to 95%, more preferably 1 to 85%, for example 5 to 60%, of a compound of formula (I). The composition is generally used for the control of pests such that a compound of formula (I) is applied at a rate of from 0.1g to 10kg per hectare, preferably from 1g to 6kg per hectare, more preferably from 1g to 1kg per hectare.

When used in a seed dressing, a compound of formula (I) is used at a rate of 0.0001g  
30 to 10g (for example 0.001g or 0.05g), preferably 0.005g to 10g, more preferably 0.005g to 4g, per kilogram of seed.

In another aspect the present invention provides an insecticidal, acaricidal, nematocidal or molluscicidal composition comprising an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor. The composition is preferably an insecticidal, acaricidal, nematocidal or molluscicidal composition.

In a still further aspect the invention provides a method of combating and controlling pests at a locus which comprises treating the pests or the locus of the pests with an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a composition comprising a compound of formula (I). The compounds of formula (I) are preferably used against insects, acarines or nematodes.

The compositions can be chosen from a number of formulation types, including dustable powders (DP), soluble powders (SP), water soluble granules (SG), water dispersible granules (WG), wettable powders (WP), granules (GR) (slow or fast release), soluble concentrates (SL), oil miscible liquids (OL), ultra low volume liquids (UL), emulsifiable concentrates (EC), dispersible concentrates (DC), emulsions (both oil in water (EW) and water in oil (EO)), micro-emulsions (ME), suspension concentrates (SC), aerosols, fogging/smoke formulations, capsule suspensions (CS) and seed treatment formulations. The formulation type chosen in any instance will depend upon the particular purpose envisaged and the physical, chemical and biological properties of the compound of formula (I).

Dustable powders (DP) may be prepared by mixing a compound of formula (I) with one or more solid diluents (for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulphur, lime, flours, talc and other organic and inorganic solid carriers) and mechanically grinding the mixture to a fine powder.

Soluble powders (SP) may be prepared by mixing a compound of formula (I) with one or more water-soluble inorganic salts (such as sodium bicarbonate, sodium carbonate or magnesium sulphate) or one or more water-soluble organic solids (such as a polysaccharide) and, optionally, one or more wetting agents, one or more dispersing agents or a mixture of said agents to improve water dispersibility/solubility. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water soluble granules (SG).

Wettable powders (WP) may be prepared by mixing a compound of formula (I) with one or more solid diluents or carriers, one or more wetting agents and, preferably, one or

more dispersing agents and, optionally, one or more suspending agents to facilitate the dispersion in liquids. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water dispersible granules (WG).

Granules (GR) may be formed either by granulating a mixture of a compound of formula (I) and one or more powdered solid diluents or carriers, or from pre-formed blank granules by absorbing a compound of formula (I) (or a solution thereof, in a suitable agent) in a porous granular material (such as pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths or ground corn cobs) or by adsorbing a compound of formula (I) (or a solution thereof, in a suitable agent) on to a hard core material (such as sands, silicates, mineral carbonates, sulphates or phosphates) and drying if necessary. Agents which are commonly used to aid absorption or adsorption include solvents (such as aliphatic and aromatic petroleum solvents, alcohols, ethers, ketones and esters) and sticking agents (such as polyvinyl acetates, polyvinyl alcohols, dextrans, sugars and vegetable oils). One or more other additives may also be included in granules (for example an emulsifying agent, wetting agent or dispersing agent).

Dispersible Concentrates (DC) may be prepared by dissolving a compound of formula (I) in water or an organic solvent, such as a ketone, alcohol or glycol ether. These solutions may contain a surface active agent (for example to improve water dilution or prevent crystallisation in a spray tank).

Emulsifiable concentrates (EC) or oil-in-water emulsions (EW) may be prepared by dissolving a compound of formula (I) in an organic solvent (optionally containing one or more wetting agents, one or more emulsifying agents or a mixture of said agents). Suitable organic solvents for use in ECs include aromatic hydrocarbons (such as alkylbenzenes or alkylnaphthalenes, exemplified by SOLVESSO 100, SOLVESSO 150 and SOLVESSO 200; SOLVESSO is a Registered Trade Mark), ketones (such as cyclohexanone or methylcyclohexanone) and alcohols (such as benzyl alcohol, furfuryl alcohol or butanol), N-alkylpyrrolidones (such as N-methylpyrrolidone or N-octylpyrrolidone), dimethyl amides of fatty acids (such as C<sub>8</sub>-C<sub>10</sub> fatty acid dimethylamide) and chlorinated hydrocarbons. An EC product may spontaneously emulsify on addition to water, to produce an emulsion with sufficient stability to allow spray application through appropriate equipment. Preparation of an EW involves obtaining a compound of formula (I) either as a liquid (if it is not a liquid at room temperature, it may be melted at a reasonable temperature, typically below 70°C) or in

solution (by dissolving it in an appropriate solvent) and then emulsifying the resultant liquid or solution into water containing one or more SFAs, under high shear, to produce an emulsion. Suitable solvents for use in EWs include vegetable oils, chlorinated hydrocarbons (such as chlorobenzenes), aromatic solvents (such as alkylbenzenes or alkyl naphthalenes) and other appropriate organic solvents which have a low solubility in water.

Microemulsions (ME) may be prepared by mixing water with a blend of one or more solvents with one or more SFAs, to produce spontaneously a thermodynamically stable isotropic liquid formulation. A compound of formula (I) is present initially in either the water or the solvent/SFA blend. Suitable solvents for use in MEs include those hereinbefore described for use in ECs or in EWs. An ME may be either an oil-in-water or a water-in-oil system (which system is present may be determined by conductivity measurements) and may be suitable for mixing water-soluble and oil-soluble pesticides in the same formulation. An ME is suitable for dilution into water, either remaining as a microemulsion or forming a conventional oil-in-water emulsion.

Suspension concentrates (SC) may comprise aqueous or non-aqueous suspensions of finely divided insoluble solid particles of a compound of formula (I). SCs may be prepared by ball or bead milling the solid compound of formula (I) in a suitable medium, optionally with one or more dispersing agents, to produce a fine particle suspension of the compound. One or more wetting agents may be included in the composition and a suspending agent may be included to reduce the rate at which the particles settle. Alternatively, a compound of formula (I) may be dry milled and added to water, containing agents hereinbefore described, to produce the desired end product.

Aerosol formulations comprise a compound of formula (I) and a suitable propellant (for example *n*-butane). A compound of formula (I) may also be dissolved or dispersed in a suitable medium (for example water or a water miscible liquid, such as *n*-propanol) to provide compositions for use in non-pressurised, hand-actuated spray pumps.

A compound of formula (I) may be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating, in an enclosed space, a smoke containing the compound.

Capsule suspensions (CS) may be prepared in a manner similar to the preparation of EW formulations but with an additional polymerisation stage such that an aqueous dispersion of oil droplets is obtained, in which each oil droplet is encapsulated by a polymeric shell and

contains a compound of formula (I) and, optionally, a carrier or diluent therefor. The polymeric shell may be produced by either an interfacial polycondensation reaction or by a coacervation procedure. The compositions may provide for controlled release of the compound of formula (I) and they may be used for seed treatment. A compound of formula (I) may also be formulated in a biodegradable polymeric matrix to provide a slow, controlled release of the compound.

A composition may include one or more additives to improve the biological performance of the composition (for example by improving wetting, retention or distribution on surfaces; resistance to rain on treated surfaces; or uptake or mobility of a compound of formula (I)). Such additives include surface active agents, spray additives based on oils, for example certain mineral oils or natural plant oils (such as soy bean and rape seed oil), and blends of these with other bio-enhancing adjuvants (ingredients which may aid or modify the action of a compound of formula (I)).

A compound of formula (I) may also be formulated for use as a seed treatment, for example as a powder composition, including a powder for dry seed treatment (DS), a water soluble powder (SS) or a water dispersible powder for slurry treatment (WS), or as a liquid composition, including a flowable concentrate (FS), a solution (LS) or a capsule suspension (CS). The preparations of DS, SS, WS, FS and LS compositions are very similar to those of, respectively, DP, SP, WP, SC and DC compositions described above. Compositions for treating seed may include an agent for assisting the adhesion of the composition to the seed (for example a mineral oil or a film-forming barrier).

Wetting agents, dispersing agents and emulsifying agents may be surface SFAs of the cationic, anionic, amphoteric or non-ionic type.

Suitable SFAs of the cationic type include quaternary ammonium compounds (for example cetyltrimethyl ammonium bromide), imidazolines and amine salts.

Suitable anionic SFAs include alkali metals salts of fatty acids, salts of aliphatic monoesters of sulphuric acid (for example sodium lauryl sulphate), salts of sulphonated aromatic compounds (for example sodium dodecylbenzenesulphonate, calcium dodecylbenzenesulphonate, butylnaphthalene sulphonate and mixtures of sodium di-*isopropyl*- and tri-*isopropyl*-naphthalene sulphonates), ether sulphates, alcohol ether sulphates (for example sodium laureth-3-sulphate), ether carboxylates (for example sodium laureth-3-carboxylate), phosphate esters (products from the reaction between one or more

fatty alcohols and phosphoric acid (predominately mono-esters) or phosphorus pentoxide (predominately di-esters), for example the reaction between lauryl alcohol and tetraphosphoric acid; additionally these products may be ethoxylated), sulphosuccinamates, paraffin or olefine sulphonates, taurates and lignosulphonates.

5           Suitable SFAs of the amphoteric type include betaines, propionates and glycines.

          Suitable SFAs of the non-ionic type include condensation products of alkylene oxides, such as ethylene oxide, propylene oxide, butylene oxide or mixtures thereof, with fatty alcohols (such as oleyl alcohol or cetyl alcohol) or with alkylphenols (such as octylphenol, nonylphenol or octylcresol); partial esters derived from long chain fatty acids or hexitol  
10   anhydrides; condensation products of said partial esters with ethylene oxide; block polymers (comprising ethylene oxide and propylene oxide); alkanolamides; simple esters (for example fatty acid polyethylene glycol esters); amine oxides (for example lauryl dimethyl amine oxide); and lecithins.

          Suitable suspending agents include hydrophilic colloids (such as polysaccharides, polyvinylpyrrolidone or sodium carboxymethylcellulose) and swelling clays (such as  
15   bentonite or attapulgate).

          A compound of formula (I) may be applied by any of the known means of applying pesticidal compounds. For example, it may be applied, formulated or unformulated, to the pests or to a locus of the pests (such as a habitat of the pests, or a growing plant liable to  
20   infestation by the pests) or to any part of the plant, including the foliage, stems, branches or roots, to the seed before it is planted or to other media in which plants are growing or are to be planted (such as soil surrounding the roots, the soil generally, paddy water or hydroponic culture systems), directly or it may be sprayed on, dusted on, applied by dipping, applied as a cream or paste formulation, applied as a vapour or applied through distribution or  
25   incorporation of a composition (such as a granular composition or a composition packed in a water-soluble bag) in soil or an aqueous environment.

          A compound of formula (I) may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods, or applied by land or aerial irrigation systems.

30           Compositions for use as aqueous preparations (aqueous solutions or dispersions) are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being added to water before use. These concentrates, which may

include DCs, SCs, ECs, EWs, MEs SGs, SPs, WPs, WGs and CSs, are often required to withstand storage for prolonged periods and, after such storage, to be capable of addition to water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Such aqueous preparations may  
5 contain varying amounts of a compound of formula (I) (for example 0.0001 to 10%, by weight) depending upon the purpose for which they are to be used.

A compound of formula (I) may be used in mixtures with fertilisers (for example nitrogen-, potassium- or phosphorus-containing fertilisers). Suitable formulation types include granules of fertiliser. The mixtures suitably contain up to 25% by weight of the  
10 compound of formula (I).

The invention therefore also provides a fertiliser composition comprising a fertiliser and a compound of formula (I).

The compositions of this invention may contain other compounds having biological activity, for example micronutrients or compounds having fungicidal activity or which  
15 possess plant growth regulating, herbicidal, insecticidal, nematocidal or acaricidal activity.

The compound of formula (I) may be the sole active ingredient of the composition or it may be admixed with one or more additional active ingredients such as a pesticide, fungicide, synergist, herbicide or plant growth regulator where appropriate. An additional active ingredient may: provide a composition having a broader spectrum of activity or  
20 increased persistence at a locus; synergise the activity or complement the activity (for example by increasing the speed of effect or overcoming repellency) of the compound of formula (I); or help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient will depend upon the intended utility of the composition. Examples of suitable pesticides include the following:

- 25 a) Pyrethroids, such as permethrin, cypermethrin, fenvalerate, esfenvalerate, deltamethrin, cyhalothrin (in particular lambda-cyhalothrin), bifenthrin, fenpropathrin, cyfluthrin, tefluthrin, fish safe pyrethroids (for example ethofenprox), natural pyrethrin, tetramethrin, s-bioallethrin, fenfluthrin, prallethrin or 5-benzyl-3-furylmethyl-(E)-(1R,3S)-2,2-dimethyl-3-(2-oxothiolan-3-ylidenemethyl)cyclopropane carboxylate;
- 30 b) Organophosphates, such as, profenofos, sulprofos, acephate, methyl parathion, azinphos-methyl, demeton-s-methyl, heptenophos, thiometon, fenamiphos, monocrotophos, profenofos, triazophos, methamidophos, dimethoate, phosphamidon, malathion, chlorpyrifos,



phosalone, terbufos, fensulfothion, fonofos, phorate, phoxim, pirimiphos-methyl, pirimiphos-ethyl, fenitrothion, fosthiazate or diazinon;

c) Carbamates (including aryl carbamates), such as pirimicarb, triazamate, cloethocarb, carbofuran, furathiocarb, ethiofencarb, aldicarb, thiofurox, carbosulfan, bendiocarb,

5 fenobucarb, propoxur, methomyl or oxamyl;

d) Benzoyl ureas, such as diflubenzuron, triflumuron, hexaflumuron, flufenoxuron or chlorfluazuron;

e) Organic tin compounds, such as cyhexatin, fenbutatin oxide or azocyclotin;

f) Pyrazoles, such as tebufenpyrad and fenpyroximate;

10 g) Macrolides, such as avermectins or milbemycins, for example abamectin, emamectin benzoate, ivermectin, milbemycin, spinosad or azadirachtin;

h) Hormones or pheromones;

i) Organochlorine compounds such as endosulfan, benzene hexachloride, DDT, chlordane or dieldrin;

15 j) Amidines, such as chlordimeform or amitraz;

k) Fumigant agents, such as chloropicrin, dichloropropane, methyl bromide or metam;

l) Chloronicotinyl compounds such as imidacloprid, thiocloprid, acetamiprid, nitenpyram or thiamethoxam;

m) Diacylhydrazines, such as tebufenozide, chromafenozide or methoxyfenozide;

20 n) Diphenyl ethers, such as diofenolan or pyriproxifen;

o) Indoxacarb;

p) Chlorfenapyr; or

q) Pymetrozine.

In addition to the major chemical classes of pesticide listed above, other pesticides  
25 having particular targets may be employed in the composition, if appropriate for the intended utility of the composition. For instance, selective insecticides for particular crops, for example stemborer specific insecticides (such as cartap) or hopper specific insecticides (such as buprofezin) for use in rice may be employed. Alternatively insecticides or acaricides specific for particular insect species/stages may also be included in the compositions (for  
30 example acaricidal ovo-larvicides, such as clofentezine, flubenzimine, hexythiazox or tetradifon; acaricidal motilicides, such as dicofol or propargite; acaricides, such as

bromopropylate or chlorobenzilate; or growth regulators, such as hydramethylnon, cyromazine, methoprene, chlorfluazuron or diflubenzuron).

Examples of fungicidal compounds which may be included in the composition of the invention are (*E*)-*N*-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide (SSF-129), 4-bromo-2-cyano-*N,N*-dimethyl-6-trifluoromethylbenzimidazole-1-sulphonamide,  $\alpha$ -[*N*-(3-chloro-2,6-xylyl)-2-methoxyacetamido]- $\gamma$ -butyrolactone, 4-chloro-2-cyano-*N,N*-dimethyl-5-*p*-tolylimidazole-1-sulfonamide (IKF-916, cyamidazosulfamid), 3-5-dichloro-*N*-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281, zoxamide), *N*-allyl-4,5,-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON65500), *N*-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy)propionamide (AC382042), *N*-(2-methoxy-5-pyridyl)-cyclopropane carboxamide, acibenzolar (CGA245704), alanycarb, aldimorph, anilazine, azaconazole, azoxystrobin, benalaxyl, benomyl, biloxazol, bitertanol, blasticidin S, bromuconazole, bupirimate, captafol, captan, carbendazim, carbendazim chlorhydrate, carboxin, carpropamid, carvone, CGA41396, CGA41397, chinomethionate, chlorothalonil, chlorozolate, clozylacon, copper containing compounds such as copper oxychloride, copper oxyquinolate, copper sulphate, copper tallate and Bordeaux mixture, cymoxanil, cyproconazole, cyprodinil, debacarb, di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat, diflumetorim, *O,O*-di-*iso*-propyl-*S*-benzyl thiophosphate, dimefluazole, dimetconazole, dimethomorph, dimethirimol, diniconazole, dinocap, dithianon, dodecyl dimethyl ammonium chloride, dodemorph, dodine, doguadine, edifenphos, epoxiconazole, ethirimol, ethyl(*Z*)-*N*-benzyl-*N*([methyl(methyl-thioethylideneaminooxycarbonyl)amino]thio)- $\beta$ -alaninate, etridiazole, famoxadone, fenamidone (RPA407213), fenarimol, fenbuconazole, fenfuram, fenhexamid (KBR2738), fenciclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, flumetover, fluoroimide, fluquinconazole, flusilazole, flutolanil, flutriafol, folpet, fuberidazole, furalaxyl, furametpyr, guazatine, hexaconazole, hydroxyisoxazole, hymexazole, imazalil, imibenconazole, iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb (SZX0722), isopropanyl butyl carbamate, isoprothiolane, kasugamycin, kresoxim-methyl, LY186054, LY211795, LY248908, mancozeb, maneb, mefenoxam, mepanipyrim, mepronil, metalaxyl, metconazole, metiram, metiram-zinc, metominostrobin, myclobutanil, neoasozin, nickel dimethyldithiocarbamate, nitrothal-*iso*propyl, nuarimol, ofurace, organomercury

compounds, oxadixyl, oxasulfuron, oxolinic acid, oxpoconazole, oxycarboxin, pefurazoate, penconazole, pencycuron, phenazin oxide, phosetyl-Al, phosphorus acids, phthalide, picoxystrobin (ZA1963), polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propiconazole, propineb, propionic acid, pyrazophos, pyrifenox, pyrimethanil, 5 pyroquilon, pyroxyfur, pyrrolnitrin, quaternary ammonium compounds, quinomethionate, quinoxifen, quintozone, sipconazole (F-155), sodium pentachlorophenate, spiroxamine, streptomycin, sulphur, tebuconazole, tecloftalam, tecnazene, tetraconazole, thiabendazole, thifluzamid, 2-(thiocyanomethylthio)benzothiazole, thiophanate-methyl, thiram, timibenconazole, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazbutyl, 10 triazoxide, tricyclazole, tridemorph, trifloxystrobin (CGA279202), triforine, triflumizole, triticonazole, validamycin A, vapam, vinclozolin, zineb and ziram.

The compounds of formula (I) may be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Examples of suitable synergists for use in the compositions include piperonyl 15 butoxide, sesamex, safroxan and dodecyl imidazole.

Suitable herbicides and plant-growth regulators for inclusion in the compositions will depend upon the intended target and the effect required.


An example of a rice selective herbicide which may be included is propanil. An example of a plant growth regulator for use in cotton is PIX™.

20 Some mixtures may comprise active ingredients which have significantly different physical, chemical or biological properties such that they do not easily lend themselves to the same conventional formulation type. In these circumstances other formulation types may be prepared. For example, where one active ingredient is a water insoluble solid and the other a water insoluble liquid, it may nevertheless be possible to disperse each active ingredient in 25 the same continuous aqueous phase by dispersing the solid active ingredient as a suspension (using a preparation analogous to that of an SC) but dispersing the liquid active ingredient as an emulsion (using a preparation analogous to that of an EW). The resultant composition is a suspoemulsion (SE) formulation.

The invention is illustrated by the following Examples:

EXAMPLE 1

This Example illustrates the preparation of compound XXIV.1, 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[(indan-1-one)oxime-3,4'-piperidine]



Step A :

Trifluoroacetic acid (27 ml) was added to a stirred solution of spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester (3.5 g, prepared according to WO 9736873) in anhydrous dichloromethane (90 ml). The reaction mixture was stirred at room temperature for 1 hour, the washed with saturated bicarbonate solution, dried over sodium sulphate and concentrated *in vacuo* to yield 2.3 g of spiro[indan-1-one-3,4'-piperidine] as a brown oil, which was used directly in the next step.

Step B:

Potassium carbonate (15.5 g) and 4-chlorocinnamyl chloride (2.1 g) were added to a solution of spiro[indan-1-one-3,4'-piperidine] (2.3 g) in acetonitrile (50 ml) under argon, and the yellow mixture was stirred at 70°C for 2 hours. The reaction mixture was filtered and the solvent evaporated *in vacuo*. The residue was partitioned between ethyl acetate and water, the organic layer was dried (sodium sulphate), filtered and concentrated *in vacuo*. The crude product was purified by chromatography [SiO<sub>2</sub>; ethyl acetate-cyclohexane: triethylamine (1:1:0.1)] to give 1.65 g of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] as a yellow oil. ; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.50 (m, 2H), 2.08 (m, 4H), 2.53 (s, 2H), 3.01 (d, J = 8Hz, 2H), 3.16 (d, J = 6.8Hz, 2H), 6.23 (dt, J = 16 Hz, 6.8 Hz, 1H), 6.45 (d, J = 16Hz, 1H), 7.19-7.71 (m, 8H); MS (ES<sup>+</sup>) 352/354 (M+H<sup>+</sup>).

Step C:


To a solution of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] (50 mg) in methanol (5 ml) were added sodium acetate (28 mg) and hydroxylamine hydrochloride (22 mg). The reaction mixture was refluxed for 2 hours and concentrated *in vacuo*. The residue was dissolved in dichloromethane, washed with water, dried (sodium sulphate) and concentrated *in vacuo*. Silica gel chromatography of the residue (eluent EtOH: ethyl acetate 9:1) afforded 40 mg of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-(*E*)-oxime-3,4'-piperidine] and 7 mg of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-(*Z*)-oxime-3,4'-piperidine] which were characterised by mass and NMR spectra.

1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-(*E*)-oxime-3,4'-piperidine]: M.p. 203°C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.60 (m, 2H), 2.14 (m, 4H), 2.80 (s, 2H), 3.01 (d, J = 10Hz, 2H), 3.19 (d, J = 6.4 Hz, 2H), 6.26 (dt, J = 16 Hz, 6.4 Hz, 1H), 6.45 (d, J = 16Hz, 1H), 7.19-7.33 (m, 7H); 7.57 (d, J = 7.6 Hz), 1H); MS (ES+) 367/369 (M+H<sup>+</sup>).

1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-(*Z*)-oxime-3,4'-piperidine]: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.60 (m, 2H), 2.20 (m, 4H), 2.70 (s, 2H), 3.03 (d, J = 8.7Hz, 2H), 3.22 (d, J = 6.0 Hz, 2H), 6.26 (dt, J = 16 Hz, 6.0 Hz, 1H), 6.45 (d, J = 16Hz, 1H), 7.15-7.34 (m, 7H); 8.32 (d, J = 7.7 Hz), 1H); MS (ES+) 367/369 (M+H<sup>+</sup>).

## EXAMPLE 2

This Example illustrates the preparation of compound XXIII.1, 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[(indan-1-ylidene)hydrazine-3,4'-piperidine]




To a solution of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] (Example 1, step B) (2.49 g) in ethanol (75 ml) was added hydrazine monohydrate (0.4 ml) and one drop of acetic acid. The reaction mixture was stirred at room temperature for 12 hours and concentrated *in vacuo*. The residue was dissolved in dichloromethane, washed

with saturated aqueous sodium bicarbonate, dried (sodium sulphate) and concentrated *in vacuo*. The solid residue was recrystallised from ethyl acetate to give 2.2 g (88%) of a white solid. M.p. 140-142°C; 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-(*E*)-ylidene]hydrazine-3,4'-piperidine]: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.45 (m, 2H), 2.15 (m, 4H), 2.34 (s, 2H), 3.03 (d, J = 10Hz, 2H), 3.22 (d, J = 6 Hz, 2H), 6.26 (dt, J = 16 Hz, 6 Hz, 1H), 6.45 (d, J = 16Hz, 1H), 7.05-7.37 (m, 7H); 7.57 (d, J = 7.2 Hz), 1H); MS (ES+) 366/368 (M+H<sup>+</sup>).

### EXAMPLE 3

This Example illustrates the preparation of compound III.3, 2-chloroisonicotinic acid [1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-ylidene-3,4'-piperidine]]hydrazide



To a solution of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] (example 1, step B) (20 mg) in ethanol (3 ml) was added 2-chloro-isonicotinic acid

hydrazide (15 mg) and one drop of sulphuric acid. The reaction mixture was refluxed for 12 hours and concentrated *in vacuo*. The residue was dissolved in dichloromethane, washed with saturated aqueous sodium bicarbonate, dried (sodium sulphate) and concentrated *in vacuo*. The solid residue was purified by preparative HPLC to give 22 mg of 2-chloroisonicotinic acid [1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-ylidene-3,4'-


piperidine]]hydrazide; M.p. 240°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.5 (m, 2H), 2.15 (m, 4H), 2.70 (s, 2H), 3.03 (m, 2H), 3.21 (m, 2H), 6.26 (dt, J = 15.8 Hz, 5.8 Hz, 1H), 6.45 (d, J = 15.8Hz, 1H), 7.1-7.9 (m, 10H); 8.50 (d, J = 4.7 Hz, 1H), 9.42 (br s, 1H); MS (ES+) 505/507 (M+H<sup>+</sup>).

Compounds I.3, II.3, V.3, XII.3, XXII.14, XXII.15, XXIII.2, XXIII.3, XXIII.4, XXIII.13, XXIII.14, XXIII.16, XXIII.17, XXIII.18, XXIII.19, XXIII.20, XXIII.21 and XXIII.22 were prepared according to procedures analogous to those described in Example 3.

5

#### EXAMPLE 4

This Example illustrates the preparation of compound III.49, 2-chloroisonicotinic acid [5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-ylidene-3,4'-piperidine]]hydrazide



10

##### Step A:

To a solution of 5-chloroindene (4.2 g) in tetrahydrofuran (30 ml) at 0°C was added dropwise lithium bis(trimethylsilyl)amide (1M in tetrahydrofuran, 70 ml) over 30 min. The resulting solution was stirred at 0°C for 1 hour then transferred by cannula into a solution of bis-(2-chloro-ethyl)-carbamic acid *tert*-butyl ester (6.8 g) in tetrahydrofuran at 0°C. The resulting solution was stirred at 0°C for 2 hours then at room temperature overnight. The solvent was evaporated *in vacuo* then the residue dissolved in dichloromethane, filtered over Hyflo, and concentrated *in vacuo*. Silica gel chromatography of the residue (eluent cyclohexane:ethyl acetate 8:2) afforded 7.3 g (82%) of a 1:1 regioisomeric mixture of 5-chloro-spiro[indene-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester and 6-chloro-spiro[indene-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.16 (m, 2H), 1.32 (s, 9H), 1.81 (m, 2H), 2.92 (m, 2H), 4.00 (m, 2H), 6.55 (6.56 for

20

the isomer) (d,  $J = 5.6$  Hz, 1H), 6.67 (6.72 for the isomer) (d,  $J = 5.6$  Hz, 1H), 6.98-7.12 (m, 3H); MS (ES+) 220/222 (M-isobutene-CO<sub>2</sub>+H<sup>+</sup>).

Step B:

- 5 To a solution of the regioisomeric mixture obtained in Step A (7 g) in tetrahydrofuran (90 ml) at 70°C under argon was added dropwise 9-borabicyclo[3.3.1]nonane (0.5M in tetrahydrofuran, 132 ml); the solution was heated at 70°C for 30 min and concentrated *in vacuo*. The residue was dissolved in dichloromethane (440 ml), cooled to 0°C and pyridinium chlorochromate (14.2 g) was added portionwise over 15 min. The resulting
- 10 mixture was refluxed for 30 min, cooled to room temperature, diluted with ether (150 ml) and filtered on Hyflo. Concentration *in vacuo* followed by silica gel chromatography of the residue (eluent cyclohexane : ethyl acetate 9:1) afforded 0.43 g of 6-chloro-spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester and 0.8 g of 5-chloro-spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester.
- 15 5-chloro-spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.39 (s, 9H), 1.70 (m, 2H), 1.92 (m, 2H), 2.65 (s, 2H), 2.87 (m, 2H), 4.26 (m, 2H), 7.39 (d,  $J = 8.1$  Hz, 1H), 7.45 (s, 1H), 7.71 (d,  $J = 8.1$  Hz); MS (ES+) 236 (M-isobutene-CO<sub>2</sub>+H<sup>+</sup>).
- 20 Step C: By analogy to the procedure described in Example 1, step A, 5-chloro-spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester (1 g) was converted into 5-chloro-spiro[indan-1-one-3,4'-piperidine] (0.75 g).

Step D: By analogy to the procedure described in Example 1, step B, 5-chloro-spiro[indan-1-one-3,4'-piperidine] (118 mg) was converted into 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] (70 mg). MS (ES+) 386/388 (M+H<sup>+</sup>).

25

Step E: By analogy to the procedure described in Example 3, 5-chloro-1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-one-3,4'-piperidine] (70 mg) was converted into the title


30 compound (46 mg). M. p. 212°C; MS (ES+) 540 (M+H<sup>+</sup>).



Compounds III.26, III.29, III.30, III.52, III.53, III.210, III.213, III.214, III.233, III.236 and III.237 were prepared according to procedures analogous to those described in Example 4.

#### EXAMPLE 5

- 5        This Example illustrates the preparation of compound III.1, 2-chloroisonicotinic acid [1'-(4-chlorobenzyl)spiro[indan-1-ylidene-3,4'-piperidine]]hydrazide



- Step A: To a solution of spiro[indan-1-one-3,4'-piperidine]-1'-carboxylic acid *tert*-butyl ester (1.5 g) and 2-chloro-isonicotinic acid hydrazide (858 mg) in ethanol (50 ml) was added one  
10 drop of sulphuric acid, and the mixture was refluxed for 5 hours. The solvent was evaporated then the residue was dissolved in dichloromethane, the organic layer washed with saturated aqueous sodium bicarbonate, dried (sodium sulphate) and concentrated *in vacuo*. The residue was purified by silica gel chromatography (eluent ethyl acetate: cyclohexane 6:4) to give 1.6 g (70%) of the hydrazide XXII.2 as a yellowish solid; M.p. 90-95°C. MS (ES+) 455 (M+H<sup>+</sup>),  
15 399 (M-isobutene+H<sup>+</sup>).

Step B:

- Trifluoroacetic acid (7.3 ml) was added to a stirred solution of the hydrazide obtained in Step A (1.46 g) in anhydrous dichloromethane (30 ml). The reaction was stirred at room  
temperature for 1 h. The reaction was washed with saturated bicarbonate solution, dried over  
20 sodium sulphate and concentrated *in vacuo* to yield 1.1 g (98%) of 2-chloroisonicotinic acid [spiro(indan-1-ylidene-3,4'-piperidine)]hydrazide XXII.1 as a yellow solid, which was used directly in the next step. M.p. 175-180°C. MS (ES+) 355 (M+H<sup>+</sup>).


Step C:

Diisopropylethylamine (0.07 ml) and 4-chlorobenzyl chloride (32 mg) were added to a solution of 2-chloroisonicotinic acid [spiro(indan-1-ylidene-3,4'-piperidine)]hydrazide (71 mg) in acetonitrile (3 ml) under argon, and the mixture was refluxed for 1 hour. The reaction mixture was diluted with ethyl acetate and washed with saturated bicarbonate solution. The organic layer was dried (sodium sulphate), filtered and the solvents removed *in vacuo*. The crude product was purified by column chromatography [SiO<sub>2</sub>; ethyl acetate- triethylamine (100:0.1)] to give 33 mg (35%) of the title compound as a white solid; M.p. 198-200 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.43 (m, 2H), 2.08 (m, 4H), 2.68 (s, 2H), 2.86 (m, 2H), 3.46 (s, 2H), 7.04-7.91 (m, 10H), 8.48 (d, J = 4.8 Hz, 1H), 9.44 (br s, 1H); MS (ES+) 352/354 (M+H<sup>+</sup>).

Compounds III.6, III.7, XXII.3, XXII.4, XXII.5, XXII.6, XXII.7, XXII.8, XXII.9, XXII.10, XXII.11, XXII.12 and XXII.13 were prepared according to procedures analogous to those described in Example 5.

#### EXAMPLE 6

This Example illustrates the preparation of compound XXIII.5



To a solution of 1'-[*trans*-3-(4-chlorophenyl)allyl]spiro[indan-1-ylidene-3,4'-piperidine]hydrazine (Example 2) (150 mg) in tetrahydrofuran (2 ml) was added 4-trifluoromethoxyphenyl isocyanate (38 mg) and the reaction mixture was stirred at room

temperature for 20 min. The solution was concentrated *in vacuo* to give 136 mg of a solid which was washed with cold dichloromethane to afford 70 mg (50%) of the title compound as a white solid; M.p. 225°C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) 1.63 (m, 2H), 1.95 (m, 4H), 2.61 (s, 2H), 2.97 (m, 2H), 3.11 (d, J = 6.4 Hz, 2H), 6.25 (dt, J = 16 Hz, 6.0 Hz, 1H), 6.45 (d, J = 16 Hz, 1H), 7.1-7.7 (m, 12H); 8.17 (s, 1H), 9.30 (s, 1H); MS (ES+) 569 (M+H<sup>+</sup>).

Compounds XXIII.6 to XXIII.12 inclusive, XXIII.15 and XXIII.23 were prepared according to procedures analogous to those described in Example 6.

10

#### EXAMPLE 7

This Example illustrates the pesticidal/insecticidal properties of compounds of formula (I). Test against were performed as follows:

*Spodoptera littoralis* (Egyptian cotton leafworm)

15 Cotton leaf discs were placed on agar in a 24-well microtiter plate and sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs were infested with 5 L<sub>1</sub> larvae. The samples were checked for mortality, repellent effect, feeding behaviour, and growth regulation 3 days after treatment (DAT). The following compounds gave at least 80% control of *Spodoptera littoralis*:

20 II.3, III.1, III.3, III.7, III.26, III.29, III.30, III.49, III.52, III.53, III.214, III.233, III.236, III.237, V.3, XVII.3, XXII.3, XXII.4, XXII.8, XXII.9, XXII.11, XXII.12, XXIII.1, XXIII.13, XXIII.16, XXIII.20, XXIII.21, XXIII.22.

*Heliothis virescens* (Tobacco budworm):

25 Eggs (0-24 h old) were placed in 24-well microtiter plate on artificial diet and treated with test solutions at an application rate of 200 ppm by pipetting. After an incubation period of 4 days, samples were checked for egg mortality, larval mortality, and growth regulation. The following compounds gave at least 80% control of *Heliothis virescens*:

30 II.3, III.1, III.3, III.6, III.7, III.26, III.29, III.30, III.49, III.52, III.53, III.210, III.213, III.214, III.233, III.236, III.237, V.3, XII.3, XXII.4, XXII.6, XXII.7, XXII.8, XXII.9, XXII.11, XXII.12, XXIII.2, XXIII.3, XXIII.13, XXIII.16, XXIII.19, XXIII.20, XXIII.21, XXIII.22.

*Plutella xylostella* (Diamond back moth):

24-well microtiter plate (MTP) with artificial diet was treated with test solutions at an application rate of 18.2 ppm by pipetting. After drying, the MTP's were infested with larvae (L2)(10-15 per well). After an incubation period of 5 days, samples were checked for larval mortality, antifeedant and growth regulation. The following compounds gave at least 80% control of *Plutella xylostella*:

II.3, III.3, III.6, III.7, III.26, III.29, III.30, III.49, III.52, III.53, III.210, III.213, III.214, III.233, III.236, III.237, V.3, XII.3, XXII.8, XXIII.1, XXIII.2, XXIII.3, XXIII.12, XXIII.13, XXIII.16, XXIII.19, XXIII.20, XXIII.21, XXIII.22, XXIV.1.

*Tetranychus urticae* (Two-spotted spider mite):

Bean leaf discs on agar in 24-well microtiter plates were sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs are infested with mite populations of mixed ages. 8 days later, discs are checked for egg mortality, larval mortality, and adult mortality. The following compounds gave at least 80% control of *Tetranychus urticae*: III.26, XXIII.21.


*Aedes aegypti* (Yellow fever mosquito):

10-15 *Aedes* larvae (L2) together with a nutrition mixture are placed in 96-well microtiter plates. Test solutions at an application rate of 2ppm are pipetted into the wells. 2 days later, insects were checked for mortality and growth inhibition. The following compounds gave at least 80% control of *Aedes aegypti*

II.3, III.3, III.7, III.26, III.29, III.30, III.49, III.52, III.53, III.213, III.214, III.233, III.236, V.3, XXII.12, XXIII.21, XXIII.22.

## CLAIMS

1. A compound of formula I



wherein X is O or NR<sup>11</sup> where R<sup>11</sup> is hydrogen, optionally substituted alkyl,

optionally substituted aryl or optionally substituted heteroaryl;

Y is a single bond, C=O, C=S or S(O)<sub>m</sub> where m is 0, 1 or 2;

R<sup>1</sup> is hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted alkylcarbonyl, aminocarbonyl, optionally substituted

alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally

substituted aryl, optionally substituted heteroaryl, optionally substituted alkoxy,

optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally

substituted heterocycloxy, cyano, optionally substituted alkenyl, optionally

substituted alkynyl, optionally substituted cycloalkyl, optionally substituted

cycloalkenyl, formyl, optionally substituted heterocyclyl, optionally substituted

alkylthio, NO or NR<sup>13</sup>R<sup>14</sup> where R<sup>13</sup> and R<sup>14</sup> are independently hydrogen, COR<sup>15</sup>,

optionally substituted alkyl, optionally substituted aryl, optionally substituted

heteroaryl, optionally substituted heterocyclyl or R<sup>13</sup> and R<sup>14</sup> together with the N atom

to which they are attached form a group -N=C(R<sup>16</sup>)-NR<sup>17</sup>R<sup>18</sup>; R<sup>15</sup> is H, optionally

substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally

substituted aryloxy optionally substituted heteroaryl, optionally substituted

heteroaryloxy or NR<sup>19</sup>R<sup>20</sup>; R<sup>16</sup>, R<sup>17</sup> and R<sup>18</sup> are each independently H or lower alkyl;

R<sup>19</sup> and R<sup>20</sup> are independently optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halogen, cyano, optionally substituted alkyl,

optionally substituted alkoxy or optionally substituted aryl;

each  $R^4$  is independently halogen, nitro, cyano, optionally substituted  $C_{1-8}$  alkyl, optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted  $C_{3-7}$  cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio or  $R^{21}R^{22}N$  where  $R^{21}$  and  $R^{22}$  are, independently, hydrogen,  $C_{1-8}$  alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-6}$  alkenyl,  $C_{3-6}$  alkynyl,  $C_{3-7}$  cycloalkyl( $C_{1-4}$ )alkyl,  $C_{2-6}$  haloalkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxycarbonyl or  $R^{21}$  and  $R^{22}$  together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two  $C_{1-6}$  alkyl groups, or 2 adjacent groups  $R^4$  together with the carbon atoms to which they are attached form a 4, 5, 6, or 7 membered carbocyclic or heterocyclic ring which may be optionally substituted by halogen; n is 0, 1, 2, 3 or 4;

each  $R_a$  is independently hydrogen, halogen, hydroxy, cyano, optionally substituted  $C_{1-8}$  alkyl, optionally substituted  $C_{2-6}$  alkenyl, optionally substituted  $C_{2-6}$  alkynyl, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted alkylaminocarbonyl, optionally substituted dialkylaminocarbonyl, optionally substituted  $C_{3-7}$  cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocyclyl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted heteroaryloxy, optionally substituted alkylthio, optionally substituted arylthio or  $R^{23}R^{24}N$  where  $R^{23}$  and  $R^{24}$  are, independently, hydrogen,  $C_{1-8}$  alkyl,  $C_{3-7}$  cycloalkyl,  $C_{3-6}$  alkenyl,  $C_{3-6}$  alkynyl,  $C_{3-7}$  cycloalkyl( $C_{1-4}$ )alkyl,  $C_{2-6}$  haloalkyl,  $C_{1-6}$  alkoxy( $C_{1-6}$ )alkyl,  $C_{1-6}$  alkoxycarbonyl or  $R^{23}$  and  $R^{24}$  together with the N atom to which they are attached form a five, six or seven-membered heterocyclic ring which may contain one or two further heteroatoms selected from O, N or S and which may be optionally substituted by one or two  $C_{1-6}$  alkyl groups, or two  $R_a$  groups attached to the same carbon atom are =O or two  $R_a$  groups attached to adjacent carbon atoms form a bond, or two  $R_a$  groups together with the carbon atom to which they are attached form a three- to

seven-membered ring, that may be saturated or unsaturated, and that may contain one or two hetero atoms selected from the group consisting of N, O and S, and which may be optionally substituted by one or two C<sub>1-6</sub> alkyl groups; or two R<sub>a</sub> groups together form a group -CH<sub>2</sub>-, -CH=CH- or -CH<sub>2</sub>CH<sub>2</sub>;


p is 0, 1, 2, 3, 4, 5 or 6; q is 0, 1, 2, 3, 4, 5 or 6 provided that p+q is 1, 2, 3, 4, 5 or 6; R<sup>8</sup> is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted alkoxy, optionally substituted aryloxy, optionally substituted alkoxycarbonyl, optionally substituted alkylcarbonyl or optionally substituted alkenylcarbonyl; or salts or N-oxides thereof.

2. An insecticidal acaricidal and nematocidal composition comprising an insecticidally, acaricidally or nematocidally effective amount of a compound of formula I as defined in claim 1.
3. A method of combating and controlling insects, acarines, nematodes or molluscs which comprises applying to a pest, to a locus of a pest, or to a plant susceptible to attack by a pest an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula I.

## ABSTRACT

CHEMICAL COMPOUNDS

An insecticidal compound of formula I



wherein X is O or NR<sup>11</sup> where R<sup>11</sup> is hydrogen, optionally substituted alkyl, optionally substituted aryl or optionally substituted heteroaryl;

Y is a single bond, C=O, C=S or S(O)<sub>m</sub> where m is 0, 1 or 2;

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>8</sup> and Ra are specified organic groups and p is 0, 1, 2, 3, 4, 5 or 6;

q is 0, 1, 2, 3, 4, 5 or 6 provided that p+q is 1, 2, 3, 4, 5 or 6; or salts or N-oxides thereof; compositions containing them and their using in controlling insects, acarines, nematodes or molluscs





PCT/IB2004/004108

